

Anti-Zebrafish JAG1B Antibody Picoband®

Catalog Number: AZQ90Y54

About JAG1B

Predicted to enable Notch binding activity. Acts upstream of or within several processes, including exocrine pancreas development; face morphogenesis; and inner ear development. Predicted to be located in membrane. Is expressed in several structures, including digestive system; endocrine system; nervous system; otic vesicle; and pharyngeal arch. Human ortholog(s) of this gene implicated in Alagille syndrome; Charcot-Marie-Tooth disease; congenital heart disease; multiple sclerosis; and tetralogy of Fallot. Orthologous to human JAG1 (jagged canonical Notch ligand 1).


Overview

Product Name	Anti-Zebrafish JAG1B Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish JAG1B Antibody Picoband® catalog # AZQ90Y54. Tested in WB, IHC, IF applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IF, IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q90Y54


Technical Details

Immunogen	E.coli-derived Zebrafish JAG1B recombinant protein (Position: Q853-V1213).
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish Immunofluorescence, 2 ug/ml, Zebrafish


Anti-Zebrafish JAG1B Antibody Picoband® (AZQ90Y54) Images



Western blot analysis of JAG1B using anti-JAG1B antibody (AZQ90Y54). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-JAG1B antigen affinity purified polyclonal antibody (AZQ90Y54) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for JAG1B at approximately 155 kDa. The expected band size for JAG1B is at 133 kDa.



IHC analysis of JAG1B using anti-JAG1B antibody (AZQ90Y54). JAG1B was detected in a paraffin-embedded section of zebrafish pancreas tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-JAG1B Antibody (AZQ90Y54) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IF analysis of JAG1B using anti-JAG1B antibody (AZQ90Y54). JAG1B was detected in a paraffin-embedded section of zebrafish embryo tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/mL rabbit anti-JAG1B Antibody (AZQ90Y54) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish JAG1B Antibody

For Research Use Only. Not for use in diagnostic procedures.