

Anti-Zebrafish HHEX Antibody Picoband®

Catalog Number: AZQ9IAV3

About HHEX

Hematopoietically-expressed homeobox protein HHEX is a protein that in humans is encoded by the HHEX gene. Homeobox genes are members of a family of transcription factors that regulate tissue development in many different organisms. Hromas et al. (1993) set out to identify homeobox genes that might play a role in hematopoiesis. And using somatic cell hybrid analysis, they mapped the HHEX gene to chromosome 10, where the HOX11 gene is located. Homeobox genes are involved in neoplastic transformation of both epithelial and hemopoietic tissues. The divergent homeobox gene HEX is expressed in the anterior visceral endoderm during early mouse development and in some adult tissues of endodermal origin, including liver and thyroid. D'Elia et al.'s findings suggested that regulation of HEX entry in the nucleus of thyrocytes may represent a critical step during human thyroid tumorigenesis.

Overview


Product Name	Anti-Zebrafish HHEX Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish HHEX Antibody Picoband® catalog #AZQ9IAV3. Tested in WB applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9IAV3

Technical Details

Immunogen	E.coli-derived zebrafish HHEX recombinant protein (Position: Y60-L228)
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish



Anti-Zebrafish HHEX Antibody Picoband® (AZQ91AV3) Images



Western blot analysis of HHEX using anti-HHEX antibody (AZQ91AV3). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates. Lane 2: whole female zebrafish tissue lysates. Lane 3: zebrafish embryo tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-HHEX antigen affinity purified polyclonal antibody (AZQ91AV3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for HHEX at approximately 26 kDa. The expected band size for HHEX is at 26 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish HHEX Antibody

For Research Use Only. Not for use in diagnostic procedures.