

Anti-Zebrafish TWIST1A/1B/2 Antibody Picoband®

Catalog Number: AZQ9PTE3

About TWIST1A/1B/2

Predicted to enable DNA-binding transcription factor activity, RNA polymerase II-specific and RNA polymerase II transcription regulatory region sequence-specific DNA binding activity. Acts upstream of or within pharyngeal system development and sprouting angiogenesis. Is expressed in several structures, including mesenchyme; mesoderm; nervous system; neural crest; and pharyngeal arch. Human ortholog(s) of this gene implicated in Sweeney-Cox syndrome; breast cancer (multiple); and synostosis (multiple). Orthologous to human TWIST1 (twist family bHLH transcription factor 1).


Overview

Product Name	Anti-Zebrafish TWIST1A/1B/2 Antibody Picoband®
Reactive Species	Zebrafish
Description	Boster Bio Anti-Zebrafish TWIST1A/1B/2 Antibody Picoband® catalog # AZQ9PTE3. Tested in WB, IHC applications. This antibody reacts with Zebrafish. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q9PTE3/Q568M7/Q5XJ18


Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of zebrafish TWIST1A/1B/2.
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.25-0.5 ug/ml, Zebrafish Immunohistochemistry(Paraffin-embedded Section), 2-5 ug/ml, Zebrafish


Anti-Zebrafish TWIST1A/1B/2 Antibody Picoband® (AZQ9PTE3) Images



Western blot analysis of TWIST1A/1B/2 using anti-TWIST1A/1B/2 antibody (AZQ9PTE3). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: zebrafish head tissue lysates, Lane 2: whole male zebrafish tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TWIST1A/1B/2 antigen affinity purified polyclonal antibody (AZQ9PTE3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for TWIST1A/1B/2 at approximately 28 kDa. The expected band size for TWIST1A/1B/2 is at 21 kDa.




IHC analysis of TWIST1A/1B/2 using anti-TWIST1A/1B/2 antibody (AZQ9PTE3). TWIST1A/1B/2 was detected in a paraffin-embedded section of zebrafish brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TWIST1A/1B/2 Antibody (AZQ9PTE3) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of TWIST1A/1B/2 using anti-TWIST1A/1B/2 antibody (AZQ9PTE3). TWIST1A/1B/2 was detected in a paraffin-embedded section of zebrafish spinal cord tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TWIST1A/1B/2 Antibody (AZQ9PTE3) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

IHC analysis of TWIST1A/1B/2 using anti-TWIST1A/1B/2 antibody (AZQ9PTE3). TWIST1A/1B/2 was detected in a paraffin-embedded section of zebrafish muscle tissue. Heat mediated antigen retrieval was performed in EDTA buffer



(pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TWIST1A/1B/2 Antibody (AZQ9PTE3) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of TWIST1A/1B/2 using anti-TWIST1A/1B/2 antibody (AZQ9PTE3). TWIST1A/1B/2 was detected in a paraffin-embedded section of zebrafish kidney tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-TWIST1A/1B/2 Antibody (AZQ9PTE3) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Zebrafish TWIST1A/1B/2 Antibody

For Research Use Only. Not for use in diagnostic procedures.