

## Anti-H4K5,8,12ac HIST1H4A Antibody

Catalog Number: CI1033

### About HIST1H4A

Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histone tails undergo numerous post-translational modifications, which either directly or indirectly alter chromatin structure to facilitate transcriptional activation or repression or other nuclear processes. In addition to the genetic code, combinations of the different histone modifications reveal the so-called "histone code". Histone methylation and demethylation is dynamically regulated by respectively histone methyl transferases and histone demethylases. Acetylation of histone H4 is associated with active genes.

### Overview

Product Name	Anti-H4K5,8,12ac HIST1H4A Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-H4K5,8,12ac HIST1H4A Antibody (Catalog# CI1033). Tested in ChIP, ChIP-seq, ELISA, Dot blot, WB, IF applications. This antibody reacts with Human, Mouse.
Application	ChIP, ChIP-seq, Dot blot, ELISA, IF, WB
Clonality	Polyclonal
Formulation	Affinity purified polyclonal antibody in PBS containing 0.05% azide and 0.05% ProClin 300.
Storage Instructions	Store at -20°C. For long-term storage, store at -80°C. Avoid multiple freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P62805

### Technical Details

Immunogen	This antibody is raised in rabbit against the region of histone H4 containing the acetylated lysines 5, 8 and 12 (H4K5,8,12ac), using a KLH-conjugated synthetic peptide.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot. Boster recommends high sensitivity ChIP-seq Kit (CK1001 & CK1002) for Chromatin Immunoprecipitation.
Form	Liquid
Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity purified

**Suggested Dilutions**

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.


If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:


Boster Bio's internal QC testing used:

User needs to optimize the dilution ratio for this antibody.


## Anti-H4K5,8,12ac HIST1H4A Antibody (CI1033) Images




ChIP assays were performed using human K562 cells, Anti-H4K5,8,12ac polyclonal antibody (Catalog # CI1033) and optimized PCR primer sets for qPCR. A titration of the antibody consisting of 0.2, 0.5, 1 and 2 ug per ChIP experiment was analysed. IgG (1 ug/IP) was used as negative IP control. QPCR was performed with primers for promoter of the active gene EIF4A2 and for a region 1 kb upstream of the GAPDH gene, used as positive controls, and for the inactive MYO10 gene and the Sat2 satellite repeat region used as negative controls.



ChIP was performed with 0.5 ug of Anti-H4K5,8,12ac polyclonal antibody (Catalog # CI1033) on sheared chromatin from 100,000 K562 cells. The IP DNA was subsequently analysed on an Illumina Genome Analyzer. Library preparation, cluster generation and sequencing were performed according to the manufacturer instructions. The 36 bp tags were aligned to the human genome using the ELAND algorithm. Figure 2 shows the signal distribution along the complete length of chromosome 2 (figure 2A) and a zoom in to a 600 kb region (figure 2B). Figure 2C and D show the enrichment in two genomic regions on chromosome 3 and 12, respectively, containing EIF4A2 and GAPDH positive controls.




To determine the titer of the antibody, an ELISA was performed using a serial dilution of Anti-H4K5,8,12ac polyclonal antibody (Catalog # CI1033) in antigen coated wells. The antigen used was a peptide containing the histone modification of interest. By plotting the absorbance against the antibody dilution, the titer of the antibody was estimated to be 1:14,500.




Western blot analysis of H4K5,8,12ac expression in HeLa whole cell lysates (25 ug, lane 1), histone extracts from HeLa cells (15 ug, lane 2), recombinant histone H2A (1 ug, lane 3), recombinant histone H2B (1 ug, lane 4), recombinant histone H3 (1 ug, lane 5) and recombinant histone H4 (1 ug, lane 6). H4K5,8,12ac was detected using Anti-H4K5,8,12ac polyclonal antibody (Catalog # CI1033) at 1/1000 dilution.

A Dot Blot analysis was performed to test the cross reactivity of Anti-H4K5,8,12ac polyclonal antibody (Catalog # CI1033) with peptides containing other histone modifications and the unmodified H4. One hundred to 0.2 pmol of the respective peptides were spotted on a membrane. The antibody was used at a dilution of 1:20,000.



This figure shows a high specificity of the antibody for the modification of interest.



Immunofluorescence images stained on HeLa cells: (Figure A Left) Cells stained with anti-H4K5,8,12ac polyclonal antibody (Catalog # CI1033) at 1/500 dilution. (Figure A Middle) Nuclei stained with DAPI. (Figure A Right) Merged images of two stains from the left and middle. Figure B, C, D and E: staining of the cells with the H4K5,8,12ac antibody after incubation of the antibody with 10ng/ul of the following blocking peptides: H4K5,8,12 unmodified (figure B), H4K5,8,12ac (figure C), H2A.ZK5,7,11ac (figure D) and H4K5,8,12,16ac (figure E).

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-H4K5,8,12ac HIST1H4A Antibody