

CDC14A (NM_003672) Human Over-expression Lysate

Catalog Number: LS008556

Transient overexpression lysate of CDC14 cell division cycle 14 homolog A (*S. cerevisiae*) (CDC14A), transcript variant 1

Overview

Product Name	CDC14A (NM_003672) Human Over-expression Lysate
Reactive Species	Human
Description	Transient overexpression lysate of CDC14 cell division cycle 14 homolog A (<i>S. cerevisiae</i>) (CDC14A), transcript variant 1
Expression Host	HEK293T
Tag	C-Myc/DDK
Detection Antibodies	Clone OT14C5, Anti-DDK (FLAG) monoclonal antibody (M30971)
Components	1 vial of 100 ug gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 ug empty vector transfected control cell lysate in RIPA buffer 1 vial of 250ul 2xSDS Sample Buffer (4% SDS, 125mM Tris-HCl pH6.8, 10% Glycerol, 0.002% Bromphenol blue, 100mM DTT)
Storage Instructions	The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.
Preparation	HEK293T cells in 10-cm dishes were transiently transfected with Transfection Reagent and 5ug TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit. Cell lysates were aliquoted and stored at -20°C before shipping.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



CDC14A (NM_003672) Human Over-expression Lysate

For Research Use Only. Not for use in diagnostic procedures.