

C22orf39 (NM 173793) Human Over-expression Lysate

Catalog Number: LS128977

Transient overexpression lysate of chromosome 22 open reading frame 39 (C22orf39), transcript variant 1

 Submit your review to Bio.com or submit your review of this product to Bio.com

ecei
ve a
\$20
Am
azo
n gif
tcar
d! Y
our
revi
ews
hel
p yo
ur f
ello
w sc
ient
ists
mak
e
the
righ
t de
cisi
ons.
Tha
nk
you
for
you
r co
ntri
buti
on.

Overview

Product Name	C22orf39 (NM_173793) Human Over-expression Lysate
Reactive Species	Human
Description	Transient overexpression lysate of chromosome 22 open reading frame 39 (C22orf39), transcript variant 1
Expression Host	HEK293T
Tag	C-Myc/DDK

Detection Antibodies	Clone OT14C5, Anti-DDK (FLAG) monoclonal antibody (M30971)
Components	For 100 µg, it contains two vials: 1 vial of 100 µg gene specific transient over-expression cell lysate in RIPA buffer 1 vial of 100 µg whole HEK293T cell lysate in RIPA buffer For 20 µg, it contains one vial: 1 vial of 20 µg lyophilized gene specific transient over-expression cell lysate
Storage Instructions	The lysate is shipped with dry ice. Upon receiving, store the sample at -80°C. Also after dilution, the protein sample should be aliquoted and stored at -80°C for long term storage. Avoid repeated freeze-thaw cycles. Lysate samples can be diluted with 2xSDS Sample Buffer provided. Lysate samples are stable for 12 months from the date of receipt when stored at -80°C.
Preparation	HEK293T cells in 10-cm dishes were transiently transfected with Transfection Reagent and 5µg TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs before collection. The cells were lysed in modified RIPA buffer (25mM Tris-HCl pH7.6, 150mM NaCl, 1% NP-40, 1mM EDTA, 1xProteinase inhibitor cocktail mix, 1mM PMSF and 1mM Na3VO4), and then centrifuged to clarify the lysate. Protein concentration was measured by BCA kit. Cell lysates were aliquoted and stored at -20°C before shipping.

[C22orf39 \(NM_173793\) Human Over-expression Lysate](#)

For Research Use Only. Not for use in diagnostic procedures.