

Anti-Mutant p53 Rabbit Monoclonal Antibody

Catalog Number: M00001-3

About TP53

The ion channels activated by glutamate are typically divided into two classes. Those that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR) while those activated by alpha-amino-3-hydroxy-5-methyl-4-isoxalone propionic acid (AMPA) are known as AMPA receptors (AMPA). The AMPAR are comprised of four distinct glutamate receptor subunits designated (GluR1-4) and they play key roles in virtually all excitatory neurotransmission in the brain (Keinänen et al., 1990; Hollmann and Heinemann, 1994). The GluR1 subunit is widely expressed throughout the nervous system. Phosphorylation of Ser-845 on GluR1 is thought to be mediated by PKA and phosphorylation of this site increases the conductance of the AMPAR (Roche et al., 1996; Banke et al., 2000). In addition, phosphorylation of this site has been linked to synaptic plasticity as well as learning and memory (Lee et al., 2003; Esteban et al., 2003).

Overview

Product Name	Anti-Mutant p53 Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Mutant p53 Rabbit Monoclonal Antibody catalog # M00001-3. Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.
Application	Flow Cytometry, IP, IF, ICC, WB
Clonality	Monoclonal AOA-20
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P04637

Technical Details

Immunogen	A synthesized peptide derived from human Mutant p53
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this

kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:


WB 1:500-1:2000

ICC/IF 1:50-1:200


IP 1:50

FC 1:50

Anti-Mutant p53 Rabbit Monoclonal Antibody (M00001-3) Images



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of p53 in (1) A431 cell lysate; (2) HEK293 cell lysate.

9 Publications Citing This Product

1. PubMed ID: 25395712, Li W, Wu D, Wei B, Wang S, Sun H, Li X, Zhang F, Zhang C, Xin Y. Afr J Tradit Complement Altern Med. 2014 Aug 23;11(5):99-104. Ecollection 2014. Anti-Tumor Effect Of Cactus Polysaccharides On Lung Squamous Carcinoma Cells (Sk-Mes-1).
2. PubMed ID: 26137081, Lactotransferrin expression is downregulated and affects the mitogen-activated protein kinase pathway in gastric cancer
3. PubMed ID: 27065079, TALENs-directed?knockout?of the?full-length?transcription?factor?Nrf1?that?represses?malignantbehaviour?of?human?hepatocellular?carcinoma?(HepG2)?cells

Visit bosterbio.com/anti-mutant-p53-rabbit-monoclonal-antibody-m00001-3-boster.html to see all 9 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Mutant p53 Rabbit Monoclonal Antibody