

Anti-STAT3 Rabbit Monoclonal Antibody

Catalog Number: M00007-2

About STAT3

The ion channels activated by glutamate are typically divided into two classes. Those that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR) while those activated by alpha-amino-3-hydroxy-5-methyl-4-isoxalone propionic acid (AMPA) are known as AMPA receptors (AMPA). The AMPAR are comprised of four distinct glutamate receptor subunits designated (GluR1-4) and they play key roles in virtually all excitatory neurotransmission in the brain (Keinänen et al., 1990; Hollmann and Heinemann, 1994). The GluR1 subunit is widely expressed throughout the nervous system. Phosphorylation of Ser-845 on GluR1 is thought to be mediated by PKA and phosphorylation of this site increases the conductance of the AMPAR (Roche et al., 1996; Banke et al., 2000). In addition, phosphorylation of this site has been linked to synaptic plasticity as well as learning and memory (Lee et al., 2003; Esteban et al., 2003).

Overview

Product Name	Anti-STAT3 Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-STAT3 Rabbit Monoclonal Antibody catalog # M00007-2. Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.
Application	IP, IF, IHC, ICC, WB
Clonality	Monoclonal GO-19
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P40763

Technical Details

Immunogen	A synthesized peptide derived from human STAT3
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this

kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:


WB 1:500-1:2000

IHC 1:50-1:200


ICC/IF 1:50-1:200

IP 1:50


Anti-STAT3 Rabbit Monoclonal Antibody (M00007-2) Images



Western blot analysis of STAT3 expression in A431 cell lysate.



Immunohistochemical analysis of paraffin-embedded mouse liver, using STAT3 Antibody.



Immunofluorescent analysis of HeLa cells, using STAT3 Antibody .

5 Publications Citing This Product

1. PubMed ID: 33542641, Zhang Q,Duan HX,Li RL,Sun JY,Liu J,Peng W,Wu CJ,Gao YX. Inducing Apoptosis and Suppressing Inflammatory Reactions in Synovial Fibroblasts are Two Important Ways for Guizhi-Shaoyao-Zhimu Decoction Against Rheumatoid Arthritis. J Inflamm Res.2021 Jan 26;14:
2. PubMed ID: 26959884, Prognostic role of STAT3 in solid tumors: a systematic review and meta-analysis
3. PubMed ID: 26549519, IL-6 Inhibits Starvation-induced Autophagy via the STAT3/Bcl-2 Signaling Pathway

Visit [bosterbio.com/anti-stat3-rabbit-monoclonal-antibody-m00007-2-boster.html](https://www.bosterbio.com/anti-stat3-rabbit-monoclonal-antibody-m00007-2-boster.html) to see all 5 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-STAT3 Rabbit Monoclonal Antibody