

## Anti-p16INK4a/CDKN2A Antibody (Monoclonal, 32C59)

Catalog Number: M00016-5

### About CDKN2A

p16(INK4A), also known as cyclin-dependent kinase inhibitor 2A(CDKN2A), or multiple tumor suppressor 1(MTS1). The p16 gene(CDKN2A) was mapped to 9p21. The p16 gene encodes a negative regulator of the cell cycle. CDKN2 plays an important role during tumorigenesis or tumor progression in a significant proportion of pancreatic adenocarcinomas. Germ-line mutations in the CDKN2A tumor-suppressor gene have been linked to the development of melanoma in some families with inherited melanoma.

### Overview


Product Name	Anti-p16INK4a/CDKN2A Antibody (Monoclonal, 32C59)
Reactive Species	Human
Description	Boster Bio Anti-p16INK4a/CDKN2A Antibody (Monoclonal, 32C59) catalog # M00016-5. Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.
Application	IF, IHC, ICC, WB
Clonality	Monoclonal 32C59
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P42771

### Technical Details


Immunogen	Synthetic peptide.
Form	Liquid
Concentration	500 ug/ml
Purification	Protein A affinity purified.
Suggested Dilutions	Western blot, 1:500-2000 Immunohistochemistry, 1:50-200 Immunocytochemistry/Immunofluorescence, 1:50-200



## Anti-p16INK4a/CDKN2A Antibody (Monoclonal, 32C59) (M00016-5) Images



Western blot analysis of CDKN2A using anti-CDKN2A antibody (M00016-5). Electrophoresis was performed on a 12% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human 293T whole cell lysates, Lane 2: human Hela whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CDKN2A antigen affinity purified monoclonal antibody (M00016-5) at 1:500 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for CDKN2A at approximately 18 kDa. The expected band size for CDKN2A is at 16 kDa.



IHC analysis of CDKN2A using anti-CDKN2A antibody (M00016-5). CDKN2A was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-CDKN2A Antibody (M00016-5) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

### Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-p16INK4a/CDKN2A Antibody (Monoclonal, 32C59)

For Research Use Only. Not for use in diagnostic procedures.