

Anti-MyD88 Monoclonal Antibody

Catalog Number: M00025

About MYD88

The ion channels activated by glutamate are typically divided into two classes. Those that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR) while those activated by alpha-amino-3-hydroxy-5-methyl-4-isoxalone propionic acid (AMPA) are known as AMPA receptors (AMPA). The AMPAR are comprised of four distinct glutamate receptor subunits designated (GluR1-4) and they play key roles in virtually all excitatory neurotransmission in the brain (Keinänen et al., 1990; Hollmann and Heinemann, 1994). The GluR1 subunit is widely expressed throughout the nervous system. Phosphorylation of Ser-845 on GluR1 is thought to be mediated by PKA and phosphorylation of this site increases the conductance of the AMPAR (Roche et al., 1996; Banke et al., 2000). In addition, phosphorylation of this site has been linked to synaptic plasticity as well as learning and memory (Lee et al., 2003; Esteban et al., 2003).

Overview


Product Name	Anti-MyD88 Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-MyD88 Monoclonal Antibody catalog # M00025. Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Monoclonal FED-13
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q99836

Technical Details


Immunogen	A synthesized peptide derived from human MyD88 Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in Drosophila, play a pivotal role in innate immune responses. TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- Kappa B and subsequent regulation of immune and inflammatory genes.
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.

Purification	Affinity-chromatography
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>WB 1:1000-1:2000</p> <p>IHC 1:50-1:200</p> <p>ICC/IF 1:50-1:200</p> <p>FC 1:100</p>


Anti-MyD88 Monoclonal Antibody (M00025) Images



Western blot analysis of MyD88 expression in Raji cell lysate.



Immunohistochemical analysis of paraffin-embedded human breast cancer, using MyD88 Antibody.



Immunofluorescent analysis of A549 cells, using MyD88 Antibody.

2 Publications Citing This Product

1. PubMed ID: 26468333, High glucose induces and activates Toll-like receptor 4 in endothelial cells of diabetic retinopathy
2. PubMed ID: 25299052, Toll-Like Receptor 4 Prompts Human Breast Cancer Cells Invasiveness via Lipopolysaccharide Stimulation and Is Overexpressed in Patients with Lymph Node Metastasis

Visit bosterbio.com/anti-myD88-antibody-m00025-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MyD88 Monoclonal Antibody