

Anti-MyD88 Monoclonal Antibody

Catalog Number: M00025


Overview

Product Name	Anti-MyD88 Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-MyD88 Monoclonal Antibody catalog # M00025. Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Monoclonal FED-13
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q99836


Technical Details

Immunogen	A synthesized peptide derived from human MyD88 Members of the Toll-like receptor (TLR) family, named for the closely related Toll receptor in Drosophila, play a pivotal role in innate immune responses. TLRs recognize conserved motifs found in various pathogens and mediate defense responses. Triggering of the TLR pathway leads to the activation of NF- Kappa B and subsequent regulation of immune and inflammatory genes.
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200 FC 1:20


Anti-MyD88 Monoclonal Antibody (M00025) Images



Western blot analysis of MYD88 using anti-MYD88 antibody (M00025). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human K562 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MYD88 antigen affinity purified monoclonal antibody (M00025) at 1:500 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for MYD88 at approximately 36 kDa. The expected band size for MYD88 is at 33 kDa.



Immunohistochemical analysis of paraffin-embedded human breast cancer, using MyD88 Antibody.



Immunofluorescent analysis of A549 cells, using MyD88 Antibody.

2 Publications Citing This Product

1. PubMed ID: 26468333, High glucose induces and activates Toll-like receptor 4 in endothelial cells of diabetic retinopathy
2. PubMed ID: 25299052, Toll-Like Receptor 4 Prompts Human Breast Cancer Cells Invasiveness via Lipopolysaccharide Stimulation and Is Overexpressed in Patients with Lymph Node Metastasis

Visit bosterbio.com/anti-myd88-antibody-m00025-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-MyD88 Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.