

Anti-Rb RB1 Rabbit Monoclonal Antibody

Catalog Number: M00039-2

About RB1

Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns (4,5) P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3) . PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors.

Overview

Product Name	Anti-Rb RB1 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Rb RB1 Rabbit Monoclonal Antibody catalog # M00039-2. Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse.
Application	IP, IF, IHC, ICC, WB
Clonality	Monoclonal FCD-18
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P06400

Technical Details

Immunogen	A synthesized peptide derived from human Rb
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:


WB 1:1000-1:2000

IHC 1:100-1:500


ICC/IF 1:100-1:500

IP 1:50


Anti-Rb RB1 Rabbit Monoclonal Antibody (M00039-2) Images




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.





All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.




Western blot analysis of Retinoblastoma expression in (1) Jurkat cell lysate; (2) MCF-7 cell lysate.


Immunohistochemical analysis of paraffin-embedded Rat lung, using the Antibody at 1:250 dilution.




Immunohistochemical analysis of paraffin-embedded Human tongue cancer, using the Antibody at 1:250 dilution.




Immunohistochemical analysis of paraffin-embedded Human renal cancer, using the Antibody at 1:250 dilution.




Immunohistochemical analysis of paraffin-embedded Human kidney, using the Antibody at 1:1000 dilution.




Immunohistochemical analysis of paraffin-embedded Mouse spleen, using the Antibody at 1:250 dilution.



Immunohistochemical analysis of paraffin-embedded human lung cancer, using Retinoblastoma Antibody.



Immunofluorescent analysis using the Antibody at 1:50 dilution.



Immunofluorescent analysis of SH-SY5Y cells, using Retinoblastoma Antibody.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Rb RB1 Rabbit Monoclonal Antibody