

Anti-Caspase-8 CASP8 Rabbit Monoclonal Antibody

Catalog Number: M00042


Overview

Product Name	Anti-Caspase-8 CASP8 Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Caspase-8 CASP8 Rabbit Monoclonal Antibody catalog # M00042. Tested in WB application. This antibody reacts with Human.
Application	WB
Clonality	Monoclonal EEG-3
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q14790

Technical Details

Immunogen	A synthesized peptide derived from human Caspase-8
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000

Anti-Caspase-8 CASP8 Rabbit Monoclonal Antibody (M00042) Images



Western blot analysis of CASP8 using anti-CASP8 antibody (M00042). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Jurkat whole cell lysates, Lane 2: human Caco-2 whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: human RT4 whole cell lysates, Lane 5: human A549 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CASP8 antigen affinity purified monoclonal antibody (M00042) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for CASP8 at approximately 50 kDa. The expected band size for CASP8 is at 55 kDa.

3 Publications Citing This Product

1. PubMed ID: 28032492, Croton Tiglium Extract Induces Apoptosis via Bax/Bcl-2 Pathways in Human Lung Cancer A549 Cells
2. PubMed ID: 26261530, Protective effect of RIP and c-FLIP in preventing liver cancer cell apoptosis induced by TRAIL
3. PubMed ID: 20081872, Inhibition of proliferation of prostate cancer cell line, PC-3, in vitro and in vivo using (%u2212)-gossypol

Visit bosterbio.com/anti-caspase-8-rabbit-monoclonal-antibody-m00042-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Caspase-8 CASP8 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.