BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-F2 Antibody (N-Term)

Catalog Number: M00044-1

About F2

Thrombin, which cleaves bonds after Arg and Lys, converts fibrinogen to fibrin and activates factors V, VII, VIII, XIII, and, in complex with thrombomodulin, protein C. Functions in blood homeostasis, inflammation and wound healing.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-F2 Antibody (N-Term)
Reactive Species	Human
Description	Boster Bio Anti-F2 Antibody (N-Term) (Catalog # M00044-1). Tested in WB application(s). This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P00734


Technical Details

Immunogen	This F2 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 46-79 amino acids from human F2.
Predicted Reactive Species	Xenopus
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	WB: 1:2000

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-F2 Antibody (N-Term) (M00044-1) Images



All lanes : Anti-F2 Antibody (N-Term) at 1:2000 dilutionLane 1: HL-60 whole cell lysatesLane 2: A549 whole cell lysatesLane 3: human plasma lysatesLysates/proteins at 20 µg per lane. SecondaryGoat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDaBlocking/Dilution buffer: 5% NFDM/TBST.

Anti-F2 Antibody (N-Term)

For Research Use Only. Not for use in diagnostic procedures.