

## Anti-Hu CD195 Purified CCR5 Monoclonal Antibody

Catalog Number: M00061

### About CCR5

CD195 / CCR5 (also known as CKR-5) is a receptor for inflammatory CC-chemokines (characterized by a pair of adjacent cysteine residues), such as MIP-1 alpha, MIP-1 beta, or RANTES. It is a G protein-associated seven-pass transmembrane protein expressed on resting T cells with memory/effector phenotype, monocytes, macrophages and immature dendritic cells. This chemokine receptor regulates the activation and directed migration of leukocytes. Importantly, along with CD4, CD195 / CCR5 functions as a major receptor for HIV. Their ligand is the viral glycoprotein gp120.

### Overview

Product Name	Anti-Hu CD195 Purified CCR5 Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Hu CD195 Purified CCR5 Monoclonal Antibody (Catalog# M00061). Tested in Flow Cytometry, IP, WB, ICC, ELISA application(s). This antibody reacts with Human.
Application	ELISA, Flow Cytometry, IP, ICC, WB
Clonality	Monoclonal T21/8
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage Instructions	Store at 2-8°C. Do not freeze.
Host	Mouse
Uniprot ID	P51681

### Technical Details

Immunogen	CCR5 peptide (Met1-Lys22) KLH conjugate. The mouse monoclonal antibody T21/8 recognizes an extracellular epitope on the N-terminus of CD195, an approximately 45 kDa G-protein coupled receptor 1 family protein expressed on resting T cells, monocytes, macrophages, and immature dendritic cells.
Predicted Reactive Species	Primate
Isotype	Mouse IgG1 kappa
Form	Liquid
Concentration	1 mg/ml
Purification	Purified by protein-A affinity chromatography.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this

kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blotting: Do not boil cell lysates prior to SDS-PAGE.

Flow cytometry: 1-4 µg/ml

## Anti-Hu CD195 Purified CCR5 Monoclonal Antibody (M00061) Images




Figure 1. Flow Cytometry validation of CCR5 using Anti-Hu CD195 Purified CCR5 Monoclonal Antibody (M00061).

Flow cytometry surface staining pattern of human peripheral blood stained using anti-human CD195 (T21/8) purified antibody (concentration in sample 3 ug/ml) GAM APC.




Figure 2. Flow Cytometry validation of CCR5 using Anti-Hu CD195 Purified CCR5 Monoclonal Antibody (M00061).

Separation of human CD195 positive lymphocytes (red-filled) from CD195 negative lymphocytes (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human CD195 (T21/8) purified antibody (concentration in sample 3 ug/ml) GAM APC.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Hu CD195 Purified CCR5 Monoclonal Antibody