

Anti-E Cadherin 1 CDH1 Antibody Picoband® (monoclonal, 9G2)

Catalog Number: M00063-2

About CDH1

CDH1 (Cadherin 1), also known as ECAD or UVO, is a protein that in humans is encoded by the CDH1 gene. Cadherin-1 is a classical member of the cadherin superfamily. By Southern analysis of DNA from a panel of mouse-human somatic cell hybrids, Mansouri et al. (1987, 1988) assigned the UVO gene to 16q (16p11-qter). Frebourg et al. (2006) found that in human embryos CDH1 is highly expressed at 4 and 5 weeks in the frontonasal prominence and at 6 weeks in the lateral and medial nasal prominences, and is therefore expressed during critical stages of lip and palate development. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells. Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7.

Overview


Product Name	Anti-E Cadherin 1 CDH1 Antibody Picoband® (monoclonal, 9G2)
Reactive Species	Human
Description	Boster Bio Anti-E Cadherin 1 CDH1 Antibody Picoband® (monoclonal, 9G2) catalog # M00063-2. Tested in Flow Cytometry, IF, IHC, ICC, WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Monoclonal 9G2
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P12830

Technical Details


Immunogen	E.coli-derived human E Cadherin recombinant protein (Position: A286-A703). Human E Cadherin shares 79.7% and 80.9% amino acid (aa) sequence identity with mouse and rat E Cadherin, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG1

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml Immunocytochemistry/Immunofluorescence, 2ug/ml Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells


Anti-E Cadherin 1 CDH1 Antibody Picoband® (monoclonal, 9G2) (M00063-2) Images



Western blot analysis of E Cadherin using anti-E Cadherin antibody (M00063-2). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human placenta tissue lysates, Lane 2: human A549 whole cell lysates, Lane 3: human HEK293 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-E Cadherin antigen affinity purified polyclonal antibody (Catalog # M00063-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for E Cadherin at approximately 130KD. The expected band size for E Cadherin is at 97KD.




Expression of RVG, NDV, alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal proteins in infected BGC and SGC cells. a Western blot analysis of RVG, NDV, alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal proteins. b Immunofluorescence analysis of P-ERK. c Immunofluorescence analysis of EMT protein markers E-cadherin. BGC and SGC cells were infected with either rL-RVG, NDV and PBS for 24 h. *P < 0.5, **P < 0.01.(rL-RVG vs NDV,rL-RVG vs NDV and PBS groups, respectively, Bar = 25 um) Index in PubMed under a CC BY license. PMID: 31640627




Effects of rL-RVG and ACB pretreated SGC cells on the alpha7-nAChR, MEK/ERK signaling pathway, epithelial/mesenchymal proteins and cell migration. a Western blot analysis of alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal protein marker. b Cell migration was detected by wound healing and transwell assay. c Immunofluorescence analysis of P-ERK. d Immunofluorescence analysis of EMT proteins E-cadherin in infected SGC cells. *P < 0.5, **P < 0.01.(rL-RVG + ACB vs rL-RVG, NDV + ACB vs NDV, PBS + ACB vs PBS, respectively, Bar = 25 um) Index in PubMed under a CC BY license. PMID: 31640627


Effects of rL-RVG and MLA pretreated cells on alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal proteins and migration of cells. a Western blot analysis of the alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal protein markers. b Cell migration was detected by wound healing and transwell assay. c




Immunofluorescence analysis of P-ERK. d Immunofluorescence analysis of EMT markers E-cadherin in infected SGC cells. *P < 0.5, **P < 0.01 (rL-RVG + MLA vs rL-RVG, NDV + MLA vs NDV, PBS + MLA vs PBS, Bar = 25 um) Index in PubMed under a CC BY license. PMID: 31640627



Effects of rL-RVG and si-RNA pretreated SGC cells on the alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal proteins and cell migration. a Western blot analysis of alpha7-nAChR, MEK/ERK signaling pathway and epithelial/mesenchymal proteins. b Cell migration was detected by wound healing and transwell assay. c Immunofluorescence analysis of P-ERK. d Immunofluorescence analysis of EMT proteins E-cadherin in infected SGC cells. *P < 0.5, **P < 0.01 (rL-RVG + si-RNA vs rL-RVG, NDV + si-RNA vs NDV, PBS + si-RNA vs PBS, Bar = 25 um) Index in PubMed under a CC BY license. PMID: 31640627




Effects of rL-RVG and corynoxetine pretreated SGC cells on alpha7-nAChR, ERK and epithelial/mesenchymal proteins and cell migration. a Western blot analysis of alpha7-nAChR, P-ERK/ERK and epithelial/mesenchymal markers. b Cell migration of SGC cells was detected by wound healing and transwell assay. c Immunofluorescence analysis of P-ERK in SGC cells. d Immunofluorescence analysis of EMT protein E-cadherin in infected SGC cells. *P < 0.5, **P < 0.01 (rL-RVG + corynoxetine vs rL-RVG, NDV + corynoxetine vs NDV, PBS + corynoxetine vs PBS, Bar = 25 um) Index in PubMed under a CC BY license. PMID: 31640627




IHC analysis of E Cadherin using anti-E Cadherin antibody (M00063-2). E Cadherin was detected in a paraffin-embedded section of human prostate cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-E Cadherin Antibody (M00063-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.


IHC analysis of E Cadherin using anti-E Cadherin antibody (M00063-2). E Cadherin was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then




incubated with 1ug/ml rabbit anti-E Cadherin Antibody (M00063-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of E Cadherin using anti-E Cadherin antibody (M00063-2). E Cadherin was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-E Cadherin Antibody (M00063-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of E Cadherin using anti-E Cadherin antibody (M00063-2). E Cadherin was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-E Cadherin Antibody (M00063-2) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IF analysis of COL4A1/E Cadherin using anti-COL4A1/E Cadherin antibody (PB9099/M00063-2) COL4A1/E Cadherin was detected in paraffin-embedded section of human colon cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/mL rabbit anti-COL4A1 Antibody (PB9099)/mouse anti E Cadherin Antibody(M00063-2) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit IgG (BA1127) /Cy3 conjugated Goat anti mouse IgG(BA1031), were used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.


IF analysis of E Cadherin using anti-E Cadherin antibody



(M00063-2). E Cadherin was detected in immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2ug/mL mouse anti-E Cadherin Antibody (M00063-2) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF analysis of E Cadherin using anti-E Cadherin antibody (M00063-2). E-cadherin was detected in paraffin-embedded section of human intestinal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/mL mouse anti-E Cadherin Antibody (M00063-2) overnight at 4°C. Biotin conjugated goat anti-mouse IgG (BA1001) was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using DyLight®488 Conjugated Avidin (BA1128). The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



Flow Cytometry analysis of A549 cells using anti- CA2 antibody (M00063-2). Overlay histogram showing A549 cells stained with M00063-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-CA2 Antibody (M00063-2, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample without incubation with primary antibody and secondary antibody (Red line) was used as a blank control.

18 Publications Citing This Product

1. PubMed ID: 10.4149/neo_2016_603, Silencing of ATPase family AAA domain-containing protein 2 inhibits migration and invasion of colorectal cancer cells.
2. PubMed ID: 10.1038/labinvest.2013.108, All-transretinoic acid ameliorates bleomycin-induced lung fibrosis by downregulating the TGF- beta 1/Smad3 signaling pathway in rats
3. PubMed ID: 10.1007/s12253-011-9380-0, Effects of Homeodomain Protein CDX2 Expression on the Proliferation and Migration of Lovo Colon Cancer Cells

Visit bosterbio.com/anti-e-cadherin-picoband-trade-antibody-monoclonal-m00063-2-boster.html to see all 18 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-E Cadherin 1 CDH1 Antibody (monoclonal, 9G2)

For Research Use Only. Not for use in diagnostic procedures.