

## Anti-Aromatase CYP19A1 Rabbit Monoclonal Antibody

Catalog Number: M00071

### About CYP19A1

Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns (4,5) P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3) . PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors.

### Overview


Product Name	Anti-Aromatase CYP19A1 Rabbit Monoclonal Antibody
Reactive Species	Human, Rat
Description	Boster Bio Anti-Aromatase CYP19A1 Rabbit Monoclonal Antibody catalog # M00071. Tested in WB, IP applications. This antibody reacts with Human, Rat.
Application	IP, WB
Clonality	Monoclonal AABB-3
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P11511

### Technical Details


Immunogen	A synthesized peptide derived from human Aromatase
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:  
WB 1:500-1:2000  
IP 1:50


## Anti-Aromatase CYP19A1 Rabbit Monoclonal Antibody (M00071) Images




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of Aromatase expression in rat brain lysate.

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



### 3 Publications Citing This Product

1. PubMed ID: 15661083, Extragonadal aromatization increases with time after ovariectomy in rats
2. PubMed ID: 15777765, Electroacupuncture enhances extragonadal aromatization in ovariectomized rats
3. PubMed ID: 23484106, Zhang D, Yang B, Zou W, Lu X, Xiong M, Wu L, Wang J, Gao J, Xu S, Zou T. Biomed Res Int. 2013;2013:301540. Doi: 10.1155/2013/301540. Epub 2012 Dec 26. Estradiol Synthesis And Release In Cultured Female Rat Bone Marrow Stem Cells.

Visit [bosterbio.com/anti-aromatase-rabbit-monoclonal-antibody-m00071-boster.html](http://bosterbio.com/anti-aromatase-rabbit-monoclonal-antibody-m00071-boster.html) to see all 3 publications.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Aromatase CYP19A1 Rabbit Monoclonal Antibody