

Anti-RAD51 Antibody (Monoclonal, 31R94)

Catalog Number: M00088-4

About RAD51

DNA repair protein RAD51 homolog 1, also known as RAD51A, is a human gene. The Rad51 gene, HsRAD51, is a homolog of RecA of Escherichia coli and functions in recombination and DNA repair. BRCA1 and BRCA2 proteins form a complex with Rad51, and these genes are thought to participate in a common DNA damage response pathway associated with the activation of homologous recombination and double-strand break repair. RAD51 is also found to interact with BRCA1 and BRCA2, which may be important for the cellular response to DNA damage. BRCA2 is shown to regulate both the intracellular localization and DNA-binding ability of this protein. Loss of these controls following BRCA2 inactivation may be a key event leading to genomic instability and tumorigenesis.

Overview

Product Name	Anti-RAD51 Antibody (Monoclonal, 31R94)
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-RAD51 Antibody (Monoclonal, 31R94) catalog # M00088-4. Tested in WB, IHC, IF, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal 31R94
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q06609


Technical Details

Immunogen	Synthetic peptide within Human Rad51 aa 3-48.
Form	Liquid
Concentration	500 ug/ml
Purification	Protein A affinity purified.

Suggested Dilutions

Western blot, 1:500-2000
Immunohistochemistry, 1:50-200
Immunofluorescence, 1:50-200
Immunocytochemistry/Immunofluorescence, 1:50-200
ImmunoPrecipitation, 1:50
Flow Cytometry (Fixed), 1:50-200

Anti-RAD51 Antibody (Monoclonal, 31R94) (M00088-4) Images



Western blot analysis of RAD51 using anti-RAD51 antibody (M00088-4). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human K562 whole cell lysates, Lane 2: human Jurkat whole cell lysates, Lane 3: human RT4 whole cell lysates, Lane 4: rat testis tissue lysates, Lane 5: mouse testis tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RAD51 antigen affinity purified monoclonal antibody (M00088-4) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for RAD51 at approximately 38 kDa. The expected band size for RAD51 is at 36 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RAD51 Antibody (Monoclonal, 31R94)

For Research Use Only. Not for use in diagnostic procedures.