

Anti-Tau MAPT Rabbit Monoclonal Antibody

Catalog Number: M00097-1

About MAPT

Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns (4,5) P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3) . PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors.

Overview

Product Name	Anti-Tau MAPT Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Tau MAPT Rabbit Monoclonal Antibody catalog # M00097-1. Tested in WB, IHC, IP applications. This antibody reacts with Human, Mouse, Rat.
Application	IP, IHC, WB
Clonality	Monoclonal GF-13
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P10636

Technical Details

Immunogen	A synthesized peptide derived from human Tau
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows:


Boster Bio's internal QC testing used:

WB 1:500-1:2000


IHC 1:100-1:500

IP 1:50

Anti-Tau MAPT Rabbit Monoclonal Antibody (M00097-1) Images



All lanes use the Antibody at 1:1k dilution for 1 hour at room temperature.



Western blot analysis of Tau expression in SH-SY5Y cell lysate.

7 Publications Citing This Product

1. PubMed ID: 32904483, Zhang YW, Li XQ, Tan WF, Fang B, Ma H. Postoperative 24-h Acute Sleep Deprivation Improves Learning and Memory Through Inhibition of Tau Phosphorylation in the Hippocampal Neurons of Splenectomized Rats. *Nat Sci Sleep*. 2020 Aug 24;12:603-613. doi:10.2147/NSS.S254
2. PubMed ID: 28934273, Han F1, Zhuang TT1, Chen JJ1, Zhu XL1,2, Cai YF1, Lu YP1. *PLoS One*. 2017 Sep 21;12(9):e0185102. doi: 10.1371/journal.pone.0185102. eCollection 2017. Novel derivative of Paeonol, Paeononilsilatic sodium, alleviates behavioral damage and hippocampal ...
3. PubMed ID: 28123476, The expression of insulin-like growth factor-1 in senior patients with diabetes and dementia

Visit bosterbio.com/anti-tau-rabbit-monoclonal-antibody-m00097-1-boster.html to see all 7 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Tau MAPT Rabbit Monoclonal Antibody