

## Anti-Tau Mapt Antibody (Monoclonal, 31M08)

Catalog Number: M00097-13

### About Mapt

The microtubule-associated proteins (MAPs) coassemble with tubulin into microtubules in vitro. Microtubule-associated protein tau appears to be enriched in axons. Tau are composed of 352 to 441 amino acids. The isoforms differ from each other by the presence or absence of 29-amino acid or 58-amino acid inserts located in the N terminus and a 31-amino repeat located in the C terminus. tau is important in establishing and maintaining neuronal morphology and is a major component of the neurofibrillary tangles (NFTs) characteristic of Alzheimer's brain. Microtubule-associated protein tau (MTBT1) is mapped to chromosome 17q21.

### Overview

Product Name	Anti-Tau Mapt Antibody (Monoclonal, 31M08)
Reactive Species	Bovine, Human
Description	Boster Bio Anti-Tau Mapt Antibody (Monoclonal, 31M08) catalog # M00097-13. Tested in IHC, WB applications. This antibody reacts with Bovine, Human.
Application	IHC, WB
Clonality	Monoclonal 31M08
Formulation	Mouse IgG in stabilizing components, 1.2% sodium acetate and 0.01mg NaN <sub>3</sub> . This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P19332

### Technical Details

Immunogen	Bovine microtubule-associated proteins (MAPs).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Mouse IgG1
Form	Lyophilized

Concentration	Adding 1 ml of PBS buffer will yield a concentration of 100 ug/ml.
Purification	Ascites
Suggested Dilutions	Immunohistochemistry (Paraffin-embedded Section), 1-2ug/ml, Human, bovine Western blot, 0.5-1ug/ml, Human, bovine

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Tau Mapt Antibody (Monoclonal, 31M08)

For Research Use Only. Not for use in diagnostic procedures.