

Anti-ERK1/2 Rabbit Monoclonal Antibody

Catalog Number: M00104-1


Overview

Product Name	Anti-ERK1/2 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ERK1/2 Rabbit Monoclonal Antibody catalog # M00104-1. Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IP, IF, ICC, WB
Clonality	Monoclonal DFF-13
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P27361/P28482


Technical Details

Immunogen	A synthesized peptide derived from human ERK1/2
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 ICC/IF 1:50-200 IP 1:20 FC 1:20


Anti-ERK1/2 Rabbit Monoclonal Antibody (M00104-1) Images



Western blot analysis of ERK1/2 Antibody expression in HepG2 whole cell lysates.




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.




ERK/NF-kappaB signaling is involved in Kv1.3-related M1 macrophages. A Respective western blotting image and densitometry analysis of ERK/NF-kappaB signaling after transfection with LV-Kv1.3 in THP-1 cells. Data were expressed as mean \pm SD (n = 4). B The phosphorylation signal data are depicted in the form of stacked histogram overlaid plots. The mean fluorescence intensity (MFI) of the ERK1/2 and NF-kappaB p65 phosphorylation signal was expressed as the mean \pm SD (n = 3). C Effects of MgTx, PD98059, and Licochalcone B on Kv1.3 and ERK/NF-kappaB signaling in LV-Kv1.3-transduced THP-1 cells were analyzed by western blotting. Data were expressed as mean \pm SD (n = 3). D Respective western blotting image and densitometry analysis of ERK/NF-kappaB signaling in HK-2 cells co-cultured with the supernatant from LV-Kv1.3-transduced THP-1 cells for 48 h. Data were expressed as mean \pm SD (n = 4). E Respective western blotting image and densitometry analysis of ERK/NF-kappaB signaling in mice subjected to UUO (7 days post-surgery) and IRI. Data were expressed as mean \pm SD (n = 4). Index in PubMed under a CC BY license. PMID: 40324999


Effects of knocking-down ADRB1 CaMKII on mPFC activity and subsequent signals in METH-sired male F1. a Levels of c-Fos protein. b The c-Fos immunostaining. Scale bar, 500 μ m /100 μ m. c Density of dendritic spine. Scale bar, 50 μ m /10 μ m. d Levels of beta-arrestin2 and PKA protein. e Levels of ERK1/2, p-ERK1/2, ERK1/2/p-ERK1/2, CREB, p-CREB, p-CREB/CREB and Δ FosB protein. F1-METH-Ctrl, METH-sired male F1 mice injected with Ctrl virus. F1-METH-KD, METH-sired male




F1 mice injected with KD virus. The data are presented as the Mean \pm SD. N.S., $P > 0.05$. * $P < 0.05$.



Effect of MYH7 R453C mutation on TGF-beta/Smad2/3, ERK1/2, NF-kappaB and PI3K/AKT pathways. (a) The protein expression of TGF-beta/Smad2/3 and ERK1/2 cascades were detected by western blotting. (b-d) The protein expression of TGF-beta1/2, p-Smad2/3/Smad2/3 and p-ERK1/2/ERK1/2 were quantitated using densitometry. (e) The protein expression of NF-kappaB signaling was detected by western blotting. (f) Quantitative analysis of the protein expression of p-NF-kappaB p65 and NF-kappaB p65. ** $p < 0.01$, **** $p < 0.0001$. $n = 3$ biologically independent samples. Index in PubMed under a CC BY license. PMID: 38862020





Effects of OPN and OVE on activation of MAPK signaling pathways in LPS-stimulated RAW264.7 cells. Expression levels of p-ERK, ERK, p-p38, p-38, p-JNK, and JNK were detected in the same samples for COX-2 detection after 24 h of LPS stimulation. (A) OVE treatment. (B) OPN treatment. All experiments were carried out in triplicates and data are presented as means \pm SDs; one-way ANOVA analysis was adopted for multiple comparisons; ### $P < 0.001$, ### $P < 0.0001$, compared to the untreated control group; *** $P < 0.001$ and **** $P < 0.0001$, compared to the LPS control group. Index in PubMed under a CC BY license. PMID: 39455284




Immunofluorescent analysis of HeLa cells, using ERK1/2 Antibody.


All lanes use the Antibody at 1:2W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2W dilution for 1 hour at room temperature.




IHC analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of mouse pancreas tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. HRP Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of rat kidney tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. HRP Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.


IHC analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. HRP Conjugated Goat Anti-rabbit IgG was




used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. HRP Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. HRP Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




IHC analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. HRP Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.


IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of mouse pancreas tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution




and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human prostate cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.


IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of rat colon tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C.




DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of rat kidney tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.


IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at



4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF analysis of ERK1/2 using anti-ERK1/2 antibody (M00104-1). ERK1/2 was detected in a paraffin-embedded section of human pancreatic cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 25 ug/mL rabbit anti-ERK1/2 Antibody (M00104-1) overnight at 4°C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



13 Publications Citing This Product

1. PubMed ID: 10.26355/eurrev_201812_16641, CXCL13 inhibition induce the apoptosis of MDA-MB-231 breast cancer cells through blocking CXCR5/ERK signaling pathway.
2. PubMed ID: -, Huanyue Cui,Xueying Liu,Jin Zhang,Ke Zhang,Dahong Yao,Shi Dong,Shushu Feng,Lu Yang, Yuyao Li,Hangyu Wang,Jian Huang,Jinhui Wang,"Rhodiola rosea L. Attenuates Cigarette Smoke and Lipopolysaccharide-Induced COPD in Rats via Inflammation Inhibition and Antioxidant and Antifibrosis Pathways",Evidence-Based Complementary and Alternative Medicine,vol. 2021,Article ID 6103158,18 pages,2021.<https://doi.org/10.1155/2021/6103158>
3. PubMed ID: -, Feng,S.,Wang,S.,Sun,S.,Su,H.,& Zhang,L.(2021).Effects of combination treatment with transcranial magnetic stimulation and bone marrow mesenchymal stem cell transplantation or Raf inhibition on spinal cord injury in rats.Molecular Medicine Reports,23,294.h

Visit [bosterbio.com/anti-erk1-2-rabbit-monoclonal-antibody-m00104-1-boster.html](https://www.bosterbio.com/anti-erk1-2-rabbit-monoclonal-antibody-m00104-1-boster.html) to see all 13 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your



reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-ERK1/2 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.