

Anti-PARP PARP1 Rabbit Monoclonal Antibody

Catalog Number: M00122-3

About PARP1

Phosphoinositide-3-kinase (PI3K) that phosphorylates PtdIns (Phosphatidylinositol), PtdIns4P (Phosphatidylinositol 4-phosphate) and PtdIns (4,5) P2 (Phosphatidylinositol 4,5-bisphosphate) to generate phosphatidylinositol 3,4,5-trisphosphate (PIP3). PIP3 plays a key role by recruiting PH domain-containing proteins to the membrane, including AKT1 and PDPK1, activating signaling cascades involved in cell growth, survival, proliferation, motility and morphology. Participates in cellular signaling in response to various growth factors.

Overview

Product Name	Anti-PARP PARP1 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PARP PARP1 Rabbit Monoclonal Antibody catalog # M00122-3. Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Monoclonal CFD-16
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P09874

Technical Details

Immunogen	A synthesized peptide derived from human PARP
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:


WB 1:500-1:2000

IHC 1:50-1:200


ICC/IF 1:50-1:200

FC 1:50


Anti-PARP PARP1 Rabbit Monoclonal Antibody (M00122-3) Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.





All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.


All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Western blot analysis of PARP expression in Jurkat cell lysate.



Immunohistochemical analysis of paraffin-embedded human testis, using PARP Antibody.



Immunofluorescent analysis of HeLa cells, using PARP Antibody .

2 Publications Citing This Product

1. PubMed ID: 33281970, Wang P,Wang C,Liu C. Antitumor effects of dioscin in A431 cells via adjusting ATM/p53-mediated cell apoptosis, DNA

damage and migration. Oncol Lett. 2021 Jan;21(1):59.doi:10.3892/ol.2020.12321.Epub 2020 Nov 19.PMID:33281970;PMCID:PMC7709553.

2. PubMed ID: 21151457, Li Q, Li Z, Xu Xy, Guo YI, Du F. Int J Mol Sci. 2010 Nov 16;11(11):4580-90. Doi: 10.3390/Ijms11114580. Neuroprotective Properties Of Picoside Ii In A Rat Model Of Focal Cerebral Ischemia.

Visit bosterbio.com/anti-parp-rabbit-monoclonal-antibody-m00122-3-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PARP PARP1 Rabbit Monoclonal Antibody