

Anti-CCND1 Antibody

Catalog Number: M00149

About CCND1

Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition. Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase. Hypophosphorylates RB1 in early G(1) phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals. Also substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity. Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex. Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner.


Overview

Product Name	Anti-CCND1 Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-CCND1 Antibody (Catalog # M00149). Tested in WB application(s). This antibody reacts with Human, Mouse.
Application	WB
Clonality	Monoclonal 1849CT557.93.5
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	P24385

Technical Details

Immunogen	This antibody is generated from a mouse immunized with a recombinant protein from human.
Predicted Reactive Species	Human
Isotype	IgG1,kappa
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	WB: 1:2000

Anti-CCND1 Antibody (M00149) Images



All lanes : Anti-CCND1 at dilution
Lane 1: MCF-7 whole cell lysate
Lane 2: HT-1080 whole cell lysate
Lane 3: L929 whole cell lysate
Lane 4: NIH/3T3 whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : kDa
Blocking/Dilution buffer: 5% NFD/MTBST.

14 Publications Citing This Product

1. PubMed ID: 24788349, Zhang H, Zhao B, Huang C, Meng Xm, Bian Eb, Li J. Plos One. 2014 May 2;9(5):E95520. Doi: 10.1371/Journal.Pone.0095520. Ecollection 2014. Melittin Restores Pten Expression By Down-Regulating Hdac2 In Human Hepatocellular Carcinoma Hepg2 Cells.
2. PubMed ID: 25240403, Liu J, Ma J, Wu Z, Li W, Zhang D, Han L, Wang F, Reindl Km, Wu E, Ma Q. BMC Cancer. 2014 Sep 20;14:686. Doi: 10.1186/1471-2407-14-686. Arginine Deiminase Augments The Chemosensitivity Of Argininosuccinate Synthetase-Deficient Pancreatic Cancer Cel...
3. PubMed ID: 25297608, Liu H, Yan Zq, Li B, Yin Sy, Sun Q, Kou Jj, Ye D, Ferns K, Liu Hy, Liu Sl. J Ovarian Res. 2014 Sep 5;7:87. Doi: 10.1186/S13048-014-0087-1. Reduced Expression Of Sox7 In Ovarian Cancer: A Novel Tumor Suppressor Through The Wnt/??-Catenin Signaling ...

Visit bosterbio.com/anti-ccnd1-antibody-m00149-boster.html to see all 14 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CCND1 Antibody

For Research Use Only. Not for use in diagnostic procedures.