

Anti-ICAM1/Cd54 Rabbit Monoclonal Antibody

Catalog Number: M00171-1


Overview

Product Name	Anti-ICAM1/Cd54 Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-ICAM1/Cd54 Rabbit Monoclonal Antibody catalog # M00171-1. Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Monoclonal DOD-9
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P05362


Technical Details

Immunogen	A synthesized peptide derived from human ICAM1
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200 FC 1:50


Anti-ICAM1/Cd54 Rabbit Monoclonal Antibody (M00171-1) Images



Western blot analysis of ICAM1 using anti-ICAM1 antibody (M00171-1). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HepG2 whole cell lysates, Lane 2: human SIHA whole cell lysates, Lane 3: human K562 whole cell lysates, Lane 4: human A431 whole cell lysates, Lane 5: human Romas whole cell lysates, Lane 6: human Hela whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ICAM1 antigen affinity purified monoclonal antibody (M00171-1) at 1: 500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for ICAM1 at approximately 100 kDa. The expected band size for ICAM1 is at 58 kDa.



IHC analysis of ICAM1 using anti-ICAM1 antibody (M00171-1). ICAM1 was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-ICAM1 Antibody (M00171-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of ICAM1 using anti-ICAM1 antibody (M00171-1). ICAM1 was detected in a paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-ICAM1 Antibody (M00171-1) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

4 Publications Citing This Product

1. PubMed ID: 25153881, Anti-inflammatory effect of the blueberry anthocyanins malvidin-3-glucoside and malvidin-3-galactoside in endothelial cells

2. PubMed ID: 23497622, Wang W, Zhao F, Fang Y, Li X, Shen L, Cao T, Zhu H. Crit Care. 2013 Mar 11;17(2):R44. Doi: 10.1186/Cc12558. Glycyrrhizin Protects Against Porcine Endotoxemia Through Modulation Of Systemic Inflammatory Response.

3. PubMed ID: 19598299, Yao Jh, Zhang Xs, Zheng Ss, Li Yh, Wang Lm, Wang Zz, Chu L, Hu Xw, Liu Kx, Tian Xf. World J Gastroenterol. 2009 Jul 14;15(26):3240-5. Prophylaxis With Carnosol Attenuates Liver Injury Induced By Intestinal Ischemia/Reperfusion.

Visit bosterbio.com/anti-icam1-rabbit-monoclonal-antibody-m00171-1-boster.html to see all 4 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ICAM1/Cd54 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.