

Anti-ICAM1 / CD54 Rabbit Monoclonal Antibody

Catalog Number: M00171-5


Overview

Product Name	Anti-ICAM1 / CD54 Rabbit Monoclonal Antibody
Reactive Species	Mouse
Description	Boster Bio Anti-ICAM1 / CD54 Rabbit Monoclonal Antibody catalog # M00171-5. Tested in WB, IHC applications. This antibody reacts with Mouse.
Application	IHC, WB
Clonality	Monoclonal 32116
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P05362


Technical Details

Immunogen	A synthesized peptide derived from human ICAM1 / CD54
Isotype	IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200


Anti-ICAM1 / CD54 Rabbit Monoclonal Antibody (M00171-5) Images



Western blot analysis of ICAM1 using anti-ICAM1 antibody (M00171-5). Electrophoresis was performed on a 8% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: mouse liver tissue lysates, Lane 2: mouse spleen tissue lysates, Lane 3: mouse kidney tissue lysates, Lane 4: mouse RAW264.7 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-ICAM1 antigen affinity purified monoclonal antibody (M00171-5) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for ICAM1 at approximately 100 kDa. The expected band size for ICAM1 is at 58 kDa.



IHC analysis of ICAM1 using anti-ICAM1 antibody (M00171-5). ICAM1 was detected in a paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-ICAM1 Antibody (M00171-5) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.



IHC analysis of ICAM1 using anti-ICAM1 antibody (M00171-5). ICAM1 was detected in a paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-ICAM1 Antibody (M00171-5) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ICAM1 / CD54 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.