

Anti-CD80 Monoclonal Antibody [8G12]

Catalog Number: M00196-1

About CD80

CD80, also known as B7-1, is a type I membrane protein that is a member of the immunoglobulin superfamily. Like the related protein CD86, this protein is expressed by antigen-presenting cells, and is the ligand for two proteins at the cell surface of T cells, CD28 and the cytotoxic T-lymphocyte-associated protein 4 (CTLA-4). Binding of this protein with CD28 antigen is a costimulatory signal for activation of the T-cell and induces T-cell proliferation and cytokine production. CTLA-4 binding negatively regulates T-cell activation and diminishes the immune response (1). Blocking the CTLA-4-CD80/CD86 interaction has been shown to enhance T-cell functions in acute lymphoblastomic leukemia (ALL), suggesting that this pathway may be an attractive target for future cancer immunotherapy (2).

Overview

| Product Name | Anti-CD80 Monoclonal Antibody [8G12] |
|----------------------|---|
| Reactive Species | Human |
| Description | Boster Bio Anti-CD80 Monoclonal Antibody [8G12] (Catalog # M00196-1). Tested in ELISA, IHC-P, ICC, IF, Flow Cytometry applications. This antibody reacts with Human. |
| Application | ELISA, Flow Cytometry, IF, IHC-P, ICC |
| Clonality | Monoclonal Clone: 8G12 |
| Formulation | CD80 Antibody is supplied in PBS containing 0.02% sodium azide and 50% glycerol. |
| Storage Instructions | CD80 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures. |
| Host | Mouse |
| Uniprot ID | P33681 |

Technical Details

| Immunogen | CD80 antibody was raised against the extracellular domain of human CD80. |
|----------------------------|--|
| Predicted Reactive Species | Mouse, Rat |
| Cross Reactivity | PD-L1 Antibody has no cross-reactivity to PD-L2. |
| Isotype | lgG2b |
| Form | Liquid |
| Concentration | 1 mg/mL |
| Purification | CD80 Antibody is supplied as protein A purified IgG2b. |




BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com


| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. |
|---------------------|---|
| | If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. |
| | Some PubMed article(s) citing the expression level of this target are as follows: |
| | Boster Bio's internal QC testing used: |
| | CD80 antibody can be used for immunohistochemistry starting at 2 - 5 ug/mL. For |
| | immunofluorescence start at 5 ug/mL. Flow cytometry at 10 ug/ml. |
| | Antibody validated: Immunohistochemistry in human samples; Immunocytochemistry in human samples; Immunofluorescence in human samples and Flow Cytometry in mouse samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher. |



Anti-CD80 Monoclonal Antibody [8G12] (M00196-1) Images




Immunocytochemistry of CD80 in transfected HEK293 cells with CD80 antibody at 1 ug/mL. Lower left: Immunocytochemistry in transfected HEK293 cells with control mouse IgG antibody at 1 ug/mL.



Immunofluorescence of CD80 in transfected HEK293 cells with CD80 antibody at 2 ug/mL.

Red: CD80 Antibody [8G12] (RF16041)


Blue: DAPI staining



Immunofluorescence of CD80 in human stomach carcinoma tissue with CD80 antibody at 20 $\mbox{ug/mL}.$

Red: CD80 Antibody [8G12] (RF16041)

Blue: DAPI staining




Immunofluorescence of CD80 in human tonsil tissue with CD80 antibody at 2 ug/mL.


Red: CD80 Antibody [8G12] (M00196-1)

Blue: DAPI staining


Immunohistochemistry of CD80 in human stomach carcinoma tissue with CD80 antibody at 5 ug/mL.







Immunohistochemistry of CD80 in human tonsil tissue with CD80 antibody at 5 ug/mL.



Flow cytometry analysis of CD80 overexpressing HEK293 cells using CD80 antibody and control mouse IgG antibody at 10 ug/ml. Blue: Untransfected HEK293 cells. Yellow: CD80 overexpressing HEK293 cells.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-CD80 Monoclonal Antibody [8G12]