

Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody

Catalog Number: M00199-1


Overview

Product Name	Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody catalog # M00199-1. Tested in WB, IHC, ICC/IF, Flow Cytometry applications. This antibody reacts with Human, Mouse.
Application	Flow Cytometry, IF, IHC, ICC, WB
Clonality	Monoclonal CEH-23
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P19544


Technical Details

Immunogen	A synthesized peptide derived from human Wilms Tumor Protein
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200 FC 1:20


Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody (M00199-1) Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Immunohistochemical analysis of paraffin-embedded mouse kidney, using Wilms Tumor Protein Antibody.




Construction of db/db model to evaluate changes in renal pathology lipid metabolism. Two groups of twenty-week-old mice were analyzed: db/m (n=6) and db/db (n=6). A-B. Representative PAS staining of glomeruli in each group, and the semiquantitative analysis of mesangial area ratio. (**



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

Western blot analysis of WT1 expression in K562 cell lysate.



4 Publications Citing This Product

1. PubMed ID: 10.1016/j.molmet.2020.101089, P2Y2R contributes to the development of diabetic nephropathy by inhibiting autophagy response
2. PubMed ID: 32987187, Dusabimana T, Kim SR, Park EJ, Je J, Jeong K, Yun SP, Kim HJ, Kim H, Park SW. P2Y2R contributes to the development of diabetic nephropathy by inhibiting autophagy response. Mol Metab. 2020 Dec;42:101089. doi:10.1016/j.molmet.2020.101089. Epub 2020 Sep 25. PMID:32987187; PMCID:PMC7568185.
3. PubMed ID: 33562139, Dusabimana T, Park EJ, Je J, Jeong K, Yun SP, Kim HJ, Kim H, Park SW. Geniposide Improves Diabetic Nephropathy by Enhancing ULK1-Mediated Autophagy and Reducing Oxidative Stress through AMPK Activation. Int J Mol Sci. 2021 Feb 6;22(4):1651. doi:10.3390/ijms22041651.

Visit bosterbio.com/anti-wilms-tumor-protein-rabbit-monoclonal-antibody-m00199-1-boster.html to see all 4 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Wilms Tumor Protein WT1 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.