

Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody

Catalog Number: M00210


Overview

Product Name	Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody catalog # M00210. Tested in WB, IP applications. This antibody reacts with Human, Mouse, Rat.
Application	IP, WB
Clonality	Monoclonal AAGE-22
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P11473


Technical Details

Immunogen	A synthesized peptide derived from human Vitamin D Receptor
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IP 1:50

Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody (M00210) Images



Western blot analysis of VDR using anti-VDR antibody (M00210). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Hela whole cell lysates, Lane 2: mouse kidney tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-VDR antigen affinity purified monoclonal antibody (Catalog # M00210) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for VDR at approximately 48 kDa. The expected band size for VDR is at 48 kDa.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.

1 Publications Citing This Product

1. PubMed ID: 26959817, Comparative Transcriptome Analysis of Fetal Skin Reveals Key Genes Related to Hair Follicle Morphogenesis in Cashmere Goats

Visit bosterbio.com/anti-vitamin-d-receptor-rabbit-monoclonal-antibody-m00210-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Vitamin D Receptor VDR Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.