

Anti-Aurora A/B Kinase Aurkb Monoclonal Antibody

Catalog Number: M00246-3


Overview

Product Name	Anti-Aurora A/B Kinase Aurkb Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Aurora A/B Kinase Aurkb Monoclonal Antibody catalog # M00246-3. Tested in IF, IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, IHC, WB
Clonality	Monoclonal
Formulation	Purified antibody at 1mg/mL in 50% PBS, 50% glycerol plus 5mM NaN ₃
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	O70126


Technical Details

Immunogen	Full length recombinant human Aurora A protein expressed in and purified from E. coli.
Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used WB: 1:1,000. IF/ICC or IHC: 1:500-1:1,000.</p> <p>For protocols, please visit https://www.bosterbio.com/protocol-and-troubleshooting/</p>

Anti-Aurora A/B Kinase Aurkb Monoclonal Antibody (M00246-3) Images



Western blot analysis of different cell lysates and recombinant protein solutions using mouse mAb to aurora A/B, M00246-3. Left: cells were treated with 100ng/mL of nocodazol, a microtubule depolymerizing agent which induces cells to halt at G2/M phase, for 6 hours prior western blotting: [1] protein standard, [2] HeLa, [3] canine A72 cells, [4] equine NBL6 cells, and [5] murine KR158 cells. Right: western blot of purified full length human recombinant aurora A, B, C as indicated. Bands at 46kDa and 38kDa correspond to aurora A and B proteins respectively, showing that M00246-3 antibody reacts with these to proteins but not aurora C.



HeLa cell cultures were stained with M00246-3 antibody (green). The antibody stains spindle poles and mitotic spindles at anaphase and concentrates on the midbody between the two daughter cells during telophase. It is therefore a useful marker of dividing cells. Cells were counterstained with our chicken polyclonal antibody to vimentin

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Aurora A/B Kinase Aurkb Monoclonal Antibody