

## Anti-NAK/TBK1 (N-term) Rabbit Monoclonal Antibody

Catalog Number: M00261

### About TBK1

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

### Overview

Product Name	Anti-NAK/TBK1 (N-term) Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-NAK/TBK1 (N-term) Rabbit Monoclonal Antibody catalog # M00261. Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, IHC, ICC, WB
Clonality	Monoclonal AHE-20
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9UHD2

### Technical Details

Immunogen	A synthesized peptide derived from human NAK/TBK1 (N-term)
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used:

WB 1:500-1:2000  
IHC 1:50-1:200  
ICC/IF 1:50-1:200

## Anti-NAK/TBK1 (N-term) Rabbit Monoclonal Antibody (M00261) Images





Figure 1. Western blot analysis of NAK/TBK1 using anti-NAK/TBK1 antibody (M00261).


Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,  
Lane 2: human 293T whole cell lysates,  
Lane 3: human MCF-7 whole cell lysates,  
Lane 4: human K562 whole cell lysates,  
Lane 5: rat liver tissue lysates,  
Lane 6: rat lung tissue lysates,  
Lane 7: mouse liver tissue lysates,  
Lane 8: mouse lung tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-NAK/TBK1 antigen affinity purified monoclonal antibody (Catalog # M00261) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for NAK/TBK1 at approximately 84 kDa. The expected band size for NAK/TBK1 is at 84 kDa.




Immunohistochemical analysis of paraffin-embedded Rat liver, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Rat stomach, using the Antibody at 1:50 dilution.


Immunohistochemical analysis of paraffin-embedded Human lung, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded Human colon cancer, using the Antibody at 1:100 dilution.




Immunohistochemical analysis of paraffin-embedded Mouse skin, using the Antibody at 1:50 dilution.




Immunohistochemical analysis of paraffin-embedded human testis, using NAK/TBK1 (N-term) Antibody.



Immunofluorescent analysis using the Antibody at 1:50 dilution.





Immunofluorescent analysis of MCF7 cells, using NAK/TBK1 (N-term) Antibody.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NAK/TBK1 (N-term) Rabbit Monoclonal Antibody