

## Anti-MITF Antibody

Catalog Number: M00269-3

### About MITF

Transcription factor that regulates the expression of genes with essential roles in cell differentiation, proliferation and survival. Binds to symmetrical DNA sequences (E-boxes) (5'-CACGTG-3') found in the promoters of target genes, such as BCL2 and tyrosinase (TYR). Plays an important role in melanocyte development by regulating the expression of tyrosinase (TYR) and tyrosinase-related protein 1 (TYRP1). Plays a critical role in the differentiation of various cell types, such as neural crest- derived melanocytes, mast cells, osteoclasts and optic cup-derived retinal pigment epithelium.


### Overview

Product Name	Anti-MITF Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-MITF Antibody (Catalog # M00269-3). Tested in WB application(s). This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Monoclonal 1607CT834.207.47
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	O75030


### Technical Details

Immunogen	This MITF antibody is generated from a mouse immunized with a recombinant protein of human MITF.
Predicted Reactive Species	Human, Mouse, Rat
Isotype	IgG1,k
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	IHC: 1:1000 WB: 1:2000

## Anti-MITF Antibody (M00269-3) Images



Immunohistochemical analysis of paraffin-embedded Human Melanoma section using Pink1. M00269-3 was diluted at 1:1000 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



All lanes : Anti-MITF Antibody at 1:2000 dilution  
Lane 1: A431 whole cell lysate  
Lane 2: PC-12 whole cell lysate  
Lane 3: NIH/3T3 whole cell lysate  
Lysates/proteins at 20 µg per lane.  
Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 59 kDa  
Blocking/Dilution buffer: 5% NFDN/TBST.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



### Anti-MITF Antibody

For Research Use Only. Not for use in diagnostic procedures.