

Anti-Caspase-3 CASP3 Rabbit Monoclonal Antibody

Catalog Number: M00334-5

About CASP3

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

Overview

Product Name	Anti-Caspase-3 CASP3 Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Caspase-3 CASP3 Rabbit Monoclonal Antibody catalog # M00334-5. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human.
Conjugate	FITC
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal AOE-3
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P42574

Technical Details

Immunogen	A synthesized peptide derived from human Caspase-3
Predicted Reactive Species	Human, Primate
Cross Reactivity	Detects ~20kDa. Does not cross-react with alphaB-crystallin, betaL-crystallin, ̢H- crystallin, gamma-crystallin, HSP25, HSP27 or HSP47 proteins.
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

WB 1:500-1:2000

IHC 1:50-1:200

ICC/IF 1:50-1:200

IP 1:50

FC 1:50

Anti-Caspase-3 CASP3 Rabbit Monoclonal Antibody (M00334-5) Images





Figure 1. Western blot analysis of Caspase-3 using anti-Caspase-3 antibody (M00334-5).


Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,
Lane 2: human Hela whole cell lysates,
Lane 3: human Jurkat whole cell lysates,
Lane 4: human Jurkat whole cell lysates,
Lane 5: human HepG2 whole cell lysates,
Lane 6: human 293T whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Caspase-3 antigen affinity purified monoclonal antibody (Catalog # M00334-5) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Caspase-3 at approximately 32 kDa. The expected band size for Caspase-3 is at 32 kDa.




Immunohistochemical analysis of paraffin-embedded human tonsil, using Caspase-3 Antibody.



Immunofluorescent analysis of Hela cells, using Caspase-3 Antibody .

Immunoprecipitate (IP) analysis using the Antibody at 1:50 dilution. (wb at 1:3K dilution)



46 Publications Citing This Product

1. PubMed ID: -, Ahmed S. Ahmed, JAK-1/STAT-3 pathway mediated role in aging cerebellar cortex degenerative changes of albino wistar rats, Translational Research in Anatomy, 2020, 100089, ISSN 2214-854X, <https://doi.org/10.1016/j.tria.2020.100089>.
2. PubMed ID: -, Gang Li, Jing Zhou, Mengyu Sun, Juren Cen, Jing Xu, Role of luteolin extracted from Clerodendrum cyrtophyllum Turcz leaves in protecting HepG2 cells from TBHP-induced oxidative stress and its cytotoxicity, genotoxicity, Journal of Functional Foods, Volume 74, 2
3. PubMed ID: 33044585, Yu Y, Wang Y, Fei X, Song Z, Xie F, Yang F, Liu X, Xu Z, Wang G. All-Trans Retinoic Acid Prevented Vein Grafts Stenosis by Inhibiting Rb-E2F Mediated Cell Cycle Progression and KLF5-RARalpha Interaction in Human Vein Smooth Muscle Cells. Cardiovasc Drugs Ther. 2020 Oct

Visit bosterbio.com/anti-caspase-3-rabbit-monoclonal-antibody-m00334-5-boster.html to see all 46 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Caspase-3 CASP3 Rabbit Monoclonal Antibody