

Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody

Catalog Number: M00372

About MPO

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

Overview

Product Name	Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody catalog # M00372. Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human.
Application	IF, IHC, ICC, WB
Clonality	Monoclonal AOCF-13
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P05164

Technical Details

Immunogen	A synthesized peptide derived from human Myeloperoxidase
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used:

	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200
--	--

Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody (M00372) Images




Figure 1. Western blot analysis of MPO using anti-MPO antibody (M00372).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HL-60 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MPO antigen affinity purified monoclonal antibody (Catalog # M00372) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MPO at approximately 84 kDa. The expected band size for MPO is at 84 kDa.

1 Publications Citing This Product

1. PubMed ID: 30043859, The Gastric Mucosa from Patients Infected with CagA or VacA Helicobacter pylori Has a Lower Level of Dual Oxidase-2 Expression than Uninfected or %u2026

Visit [bosterbio.com/anti-myeloperoxidase-rabbit-monoclonal-antibody-m00372-boster.html](https://www.bosterbio.com/anti-myeloperoxidase-rabbit-monoclonal-antibody-m00372-boster.html) to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Myeloperoxidase MPO Rabbit Monoclonal Antibody