BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-CCR7 Antibody (N-term)

Catalog Number: M00390-1

About CCR7

Receptor for the MIP-3-beta chemokine. Probable mediator of EBV effects on B-lymphocytes or of normal lymphocyte functions.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a $20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Overview

Product Name	Anti-CCR7 Antibody (N-term)
Reactive Species	Human
Description	Boster Bio Anti-CCR7 Antibody (N-term) (Catalog # M00390-1). Tested in WB application(s). This antibody reacts with Human.
Application	WB
Clonality	Monoclonal 1987CT875.52.22
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	P32248

Technical Details


Immunogen	This CCR7 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 26-57 amino acids from the N-terminal region of human CCR7.
Predicted Reactive Species	Human
Isotype	IgG1,kappa
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	WB: 1:4000 Tested Species: In-house tested species with positive results. Other applications have not been tested. Optimal dilutions

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

	should be determined by end users.

BOSTER BIOLOGICAL TECHNOLOGY
3942 B Valley Ave, Pleasanton, CA 94566
888-466-3604 | support@bosterbio.com | www.bosterbio.com

Anti-CCR7 Antibody (N-term) (M00390-1) Images



All lanes : Anti-CCR7 Antibody (N-term) at 1:4000 dilutionLane 1: K562 whole cell lysateLane 2: Daudi whole cell lysateLysates/proteins at 20 µg per lane. SecondaryGoat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 43 kDaBlocking/Dilution buffer: 5% NFDM/TBST.

Anti-CCR7 Antibody (N-term)

For Research Use Only. Not for use in diagnostic procedures.