

## Anti- IL 17A Monoclonal Antibody

Catalog Number: M00421-2

### About IL17a

Interleukin-17A is a protein that in humans is encoded by the IL17A gene. The protein encoded by this gene is a proinflammatory cytokine produced by activated T cells. This cytokine regulates the activities of NF-kappaB and mitogen-activated protein kinases. This cytokine can stimulate the expression of IL6 and cyclooxygenase-2 (PTGS2/COX-2), as well as enhance the production of nitric oxide (NO). High levels of this cytokine are associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis.

### Overview

Product Name	Anti- IL 17A Monoclonal Antibody
Reactive Species	Mouse
Description	Boster Bio Anti- IL 17A Monoclonal Antibody catalog # M00421-2. Tested in IHC-P applications. This antibody reacts with Mouse.
Application	IHC-P
Clonality	Monoclonal AFIF-9
Formulation	PBS, pH 7.0. Contains no stabilizers or preservatives
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q62386


### Technical Details

Immunogen	Mouse IL-17A cross-linked to OVA
Predicted Reactive Species	Primate
Recommended Detection Systems	Boster recommends HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0001-1) for IHC(P) .
Cross Reactivity	No cross reactivity with other proteins.
Isotype	Mouse IgG1, kappa
Form	Lyophilized
Concentration	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Mouse

## Anti- IL 17A Monoclonal Antibody (M00421-2) Images



IHC analysis of IL17A using anti-IL17A antibody (M00421-2). IL17A was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5ug/ml mouse anti-IL17A Antibody (M00421-2) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1021) with DAB as the chromogen.

## 1 Publications Citing This Product

1. PubMed ID: 10.1038/s41401-021-00706-4, Pirfenidone ameliorates silica-induced lung inflammation and fibrosis in mice by inhibiting the secretion of interleukin-17A

Visit [bosterbio.com/anti-il-17a-monoclonal-antibody-m00421-2-boster.html](https://bosterbio.com/anti-il-17a-monoclonal-antibody-m00421-2-boster.html) to see all 1 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti- IL 17A Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.