

Anti-SYK Purified Monoclonal Antibody

Catalog Number: M00490

About SYK

Syk is a cytoplasmic protein tyrosine kinase that translocates to the plasma membrane upon B cell antigen receptor (BCR) or the high-affinity IgE receptor (FcεRI) triggering, and phosphorylates downstream adaptor proteins, thereby providing docking sites for initiation of subsequent signaling pathways, such as calcium mobilization, cytoskeleton remodeling, or transcription of specific genes. Syk binds to the receptor assemblies through interactions of its pair of SH2 domains with ITAM motives of the receptor, which have been phosphorylated by Src-family kinases. These kinases also help to activate Syk by phosphorylation of its activation loop.

Overview

Product Name	Anti-SYK Purified Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-SYK Purified Monoclonal Antibody (Catalog# M00490). Tested in IP, WB, IHC-P, ICC application(s). This antibody reacts with Human, Rat, Mouse.
Application	IP, IHC-P, ICC, WB
Clonality	Monoclonal SYK-01
Formulation	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage Instructions	Store at 2-8°C. Do not freeze.
Host	Mouse
Uniprot ID	P43405

Technical Details

Immunogen	Recombinant fragment (aa 5-360) of human Syk. The antibody SYK-01 reacts with protein tyrosine kinase p72Syk (Syk; an intracellular antigen), which is required for the transduction of signals through the B cell antigen receptor (BCR) and the high affinity IgE receptor (FcεRI).
Predicted Reactive Species	Primate
Isotype	Mouse IgG1
Form	Liquid
Concentration	1 mg/ml
Purification	Purified by protein-A affinity chromatography.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blotting: 1-2 ug/ml.

Immunohistochemistry (paraffin sections): 5 ug/ml; positive tissue: tonsil B cells.

Anti-SYK Purified Monoclonal Antibody (M00490) Images




Figure 2. Immunocytochemistry staining of SYK using Anti-SYK Purified Monoclonal Antibody (M00490).

Immunocytochemistry analysis of Syk in human primary fibroblasts using anti-Syk (SYK-01; green). Actin cytoskeleton was decorated by phalloidin (red) and cell nuclei stained with DAPI (blue).




Figure 3. Immunocytochemistry staining of SYK using Anti-SYK Purified Monoclonal Antibody (M00490).

Immunocytochemistry analysis of Syk in human HeLa cell line using anti-Syk (SYK-01; green). Actin cytoskeleton was decorated by phalloidin (red) and cell nuclei stained with DAPI (blue).




Figure 1. Western blotting validation for Anti-SYK Purified Monoclonal Antibody M00490

Western blotting analysis (non-reducing conditions) of whole cell lysate of RAMOS human Burkitt lymphoma cell line (1)

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SYK Purified Monoclonal Antibody