

## Anti-AFP Antibody

Catalog Number: M00522-2

### About AFP

Binds copper, nickel, and fatty acids as well as, and bilirubin less well than, serum albumin. Only a small percentage (less than 2%) of the human AFP shows estrogen-binding properties.


### Overview

Product Name	Anti-AFP Antibody
Reactive Species	Human
Description	Boster Bio Anti-AFP Antibody (Catalog # M00522-2). Tested in WB, IHC-P application(s). This antibody reacts with Human.
Application	IHC-P, WB
Clonality	Monoclonal 1087CT8.3.5
Formulation	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Mouse
Uniprot ID	P02771


### Technical Details

Immunogen	This AFP antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 281-609 amino acids from human AFP.
Predicted Reactive Species	Human, Mouse
Isotype	IgG2b,k
Purification	This antibody is purified through a protein G column, followed by dialysis against PBS.
Suggested Dilutions	WB: 1:2000 IHC-P: 1:25


## Anti-AFP Antibody (M00522-2) Images



Anti-AFP Antibody at 1:500 dilution + HepG2 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa. Blocking/Dilution buffer: 5% NFDN/TBST.



Anti-AFP Antibody at 1:2000 dilution + human fetal Liver lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 68 kDa. Blocking/Dilution buffer: 5% NFDN/TBST.



M00522-2 staining AFP in human liver tissue sections by Immunohistochemistry (IHC-P -paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hours at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

### Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-AFP Antibody

For Research Use Only. Not for use in diagnostic procedures.