

Anti-Protein Wnt-3a Wnt3a Monoclonal Antibody

Catalog Number: M00636

About WNT3A

TRPC6, also known as TRP6, short transient receptor potential channel 6 and transient receptor potential cation channel subfamily C member 6, is thought to form a receptor-activated non-selective calcium permeant cation channel. TRPC6 is probably operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G-protein coupled receptors. It is activated by diacylglycerol (DAG) in a membrane-delimited fashion, independently of protein kinase C and may not to be activated by intracellular calcium store depletion. Defects in this gene are a cause of focal segmental glomerulosclerosis (FSGS). Expression of this protein has been reported in tissues such as placenta, lung, spleen, ovary, small intestine, and renal podocytes. Immunohistochemistry studies using polyclonal antibodies to this target have shown moderate to strong staining in cell types such as neurons, breast, respiratory, squamous and prostate epithelium, epidermis, placental trophoblasts, dendritic cells, and subsets of immune cells, and faint to moderate staining of adrenal, colon, ganglion cells, hepatocytes, heart, and testis.

Overview

Product Name	Anti-Protein Wnt-3a Wnt3a Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-Protein Wnt-3a Wnt3a Monoclonal Antibody catalog # M00636. Tested in ELISA, WB applications. This antibody reacts with Human.
Application	ELISA, WB
Clonality	Monoclonal 6H11-H11-G7
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P56704

Technical Details

Immunogen	Purified recombinant Wnt3a protein fragments expressed in E.coli.
Predicted Reactive Species	Chimpanzee
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml





888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB 1:500-2000 ELISA 1:10000-20000



Anti-Protein Wnt-3a Wnt3a Monoclonal Antibody (M00636) Images




Figure 1. Western blotting validation for Anti-Protein Wnt-3a Wnt3a Monoclonal Antibody M00636

Western blot detection of Wnt3a in MCF-7

1 Publications Citing This Product

1. PubMed ID: 32463570, Xu T,Pan L,Li L,Hu S,Zhou H,Yang C,Yang J,Li H,Liu Y,Meng X,Li J.MicroRNA-708 modulates Hepatic Stellate Cells activation and enhances extracellular matrix accumulation via direct targeting TMEM88. J Cell Mol Med. 2020 Jul;24(13):7127-7140.doi: 10.1111/jc

Visit bosterbio.com/anti-protein-wnt-3a-wnt3a-monoclonal-antibody-m00636-boster.html to see all 1 publications.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Protein Wnt-3a Wnt3a Monoclonal Antibody