

Anti-Cytochrome P450 2E1/CYP2E1 Antibody Picoband® (monoclonal, 2C7G1) Cy3 Conjugated

Catalog Number: M00672-2-Cy3

About CYP2E1

Cytochrome P450 2E1 (abbreviated CYP2E1), a member of the cytochrome P450 mixed-function oxidase system, is involved in the metabolism of xenobiotics in the body. In humans, the CYP2E1 enzyme is encoded by the CYP2E1 gene. It is mapped to 10q26.3. While it is involved in the oxidative metabolism of a small range of substrates (mostly small polar molecules), there are many important drug interactions mediated by CYP2E1. Most drugs undergo deactivation by CYP2E1, either directly or by facilitated excretion from the body. Also, many substances are bioactivated by CYP2E1 to form their active compounds. In addition, CYP2E1 is an important enzyme for the conversion of ethanol to acetaldehyde and to acetate in humans. In the conversion sequence of acetyl-CoA to glucose, CYP2E1 transforms acetone via acetol into propylene glycol and methylglyoxal, the precursors of pyruvate, acetate and lactate.

Overview

Product Name	Anti-Cytochrome P450 2E1/CYP2E1 Antibody Picoband® (monoclonal, 2C7G1) Cy3 Conjugated
Reactive Species	Human, Mouse, Rat
Application	Recommended applications are based on the parent unconjugated antibody (WB). Customers may select suitable applications according to their experimental needs.
Clonality	Monoclonal 2C7G1
Formulation	Each vial contains 50% glycerol, 0.9% NaCl, 0.2% Na ₂ HPO ₄ , 0.02% NaN ₃ .
Storage Instructions	At -20°C for one year from date of receipt. Avoid repeated freezing and thawing. Protect from light.
Host	Mouse
Uniprot ID	P05181

Technical Details

Immunogen	E.coli-derived human Cytochrome P450 2E1/CYP2E1 recombinant protein (Position: H355-S493).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Liquid
Concentration	0.5 mg/mL
Purification	Immunogen affinity purified.
Conjugate	Cy3

	Excitation Wavelength: 554 nm Emission Wavelength: 568 nm
Suggested Dilutions	Optimal dilutions should be determined by end users.

5 Publications Citing This Product

1. PubMed ID: 10.3109/13880209.2014.898075, Effect of ammonium pyrrolidine dithiocarbamate (PDTC) on NF-kappaB activation and CYP2E1 content of rats with immunological liver injury
2. PubMed ID: 10.1016/j.apsb.2017.04.002, Schisandra sphenanthera extract (Wuzhi Tablet) protects against chronic-binge and acute alcohol-induced liver injury by regulating the NRF2-ARE pathway in mice
3. PubMed ID: 10.1016/j.toxlet.2011.11.030, Furocoumarins affect hepatic cytochrome P450 and renal organic ion transporters in mice

Visit bosterbio.com/anti-cytochrome-p450-2e1-cyp2e1-picoband-trade-antibody-m00672-2-boster.html to see all 5 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cytochrome P450 2E1/CYP2E1 Antibody (monoclonal, 2C7G1) - Cy3

For Research Use Only. Not for use in diagnostic procedures.