

Anti-STAT5b Rabbit Monoclonal Antibody

Catalog Number: M00681


Overview

Product Name	Anti-STAT5b Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-STAT5b Rabbit Monoclonal Antibody catalog # M00681. Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.
Application	IP, IF, IHC, ICC, WB
Clonality	Monoclonal HDF-19
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P51692


Technical Details

Immunogen	A synthesized peptide derived from human STAT5b
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200 IP 1:20

Anti-STAT5b Rabbit Monoclonal Antibody (M00681) Images





Western blot analysis of STAT5B using anti-STAT5B antibody (M00681). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human 293T whole cell lysates, Lane 2: human K562 whole cell lysates, Lane 3: human Hela whole cell lysates, Lane 4: human Raji whole cell lysates, Lane 5: rat lung tissue lysates, Lane 6: rat liver tissue lysates, Lane 7: mouse lung tissue lysates, Lane 8: mouse liver tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-STAT5B antigen affinity purified monoclonal antibody (M00681) at 1:1000 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for STAT5B at approximately 90 kDa. The expected band size for STAT5B is at 90 kDa.



Western blot analysis of STAT5B using anti-STAT5B antibody (M00681). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human K562 whole cell lysates, Lane 2: human HEL whole cell lysates, Lane 3: human Jurkat whole cell lysates, Lane 4: human Hela whole cell lysates, Lane 5: rat thymus tissue lysates, Lane 6: rat skeletal muscle tissue lysates, Lane 7: mouse thymus tissue lysates, Lane 8: mouse skeletal muscle tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-STAT5B antigen affinity purified monoclonal antibody (M00681) at 1: 1000 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for STAT5B at approximately 90-95 kDa. The expected band size for STAT5B is at 90 kDa.

Immunohistochemical analysis of paraffin-embedded human breast, using STAT5b Antibody.



Immunoprecipitating (IP) STAT5B in K562 whole cell lysate. Western blot analysis of STAT5B using anti-STAT5B antibody (M00681); Lane 1: K562 whole cell lysates (30ug); Lane 2: Rabbit control IgG instead of anti-STAT5B antibody in K562 whole cell lysate; Lane 3: anti-STAT5B antibody (2ug) + K562 whole cell lysate (500ug). After electrophoresis, proteins were transferred to a membrane. Then the membrane was incubated with rabbit anti-STAT5B antigen affinity purified monoclonal antibody (M00681) at a dilution of 0.5 ug/mL and probed with a goat anti-rabbit IgG-HRP secondary antibody (Catalog # BA1054). The signal is developed using ECL Plus Western Blotting Substrate (Catalog # AR1196-200). A specific band was detected for STAT5B at approximately 90 kDa. The expected band size for STAT5B is at 90 kDa.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-STAT5b Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.