

Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10)

Catalog Number: M00714

About EMD

APOBEC3G (apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 3G) is a human enzyme encoded by the APOBEC3G gene. This gene is a member of the cytidine deaminase gene family. It is one of seven related genes or pseudogenes found in a cluster, thought to result from gene duplication, on chromosome 22. Members of the cluster encode proteins that are structurally and functionally related to the C to U RNA-editing cytidine deaminase APOBEC1. It is thought that the proteins may be RNA editing enzymes and have roles in growth or cell cycle control. The protein encoded by this gene has been found to be a specific inhibitor of human immunodeficiency virus-1 (HIV-1) infectivity.

Overview

Product Name	Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10)
Reactive Species	Human
Description	Boster Bio Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) catalog # M00714. Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human.
Application	Flow Cytometry, IHC, ICC, WB
Clonality	Monoclonal 5A10
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P50402

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human Emerin, different from the related mouse sequence by eight amino acids, and from the related rat sequence by nine amino acids.
Predicted Reactive Species	Hepatitis Virus
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(P), IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG1
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml</p> <p>Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml</p> <p>Immunohistochemistry (Frozen Section), 0.5-1ug/ml</p> <p>Immunocytochemistry, 0.5-1ug/ml</p> <p>Flow Cytometry, 1-3ug/1x10⁶ cells</p>

Anti-Emerin EMD Antibody Picoband™ (monoclonal, 5A10) (M00714) Images




Figure 1. Western blot analysis of Emerin using anti-Emerin antibody (M00714).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,
Lane 2: human placenta tissue lysates,
Lane 3: human Caco-2 whole cell lysates,
Lane 4: human HepG2 whole cell lysates,
Lane 5: Rabbit IgG,
Lane 6: Marker 1113,
Lane 7: human Jurkat whole cell lysates.
Lane 8: human MDA-MB-453 whole cell lysates,
Lane 9: human SK-OV-3 whole cell lysates,
Lane 10: human SW620 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Emerin affinity purified monoclonal antibody (Catalog # M00714) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.





Figure 2. IHC analysis of Emerin using anti-Emerin antibody (M00714).

Emerin was detected in paraffin-embedded section of human gastric cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml mouse anti-Emerin Antibody (M00714) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1021) with DAB as the chromogen.

Figure 3. IHC analysis of Emerin using anti-Emerin antibody (M00714).

Emerin was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml mouse anti-Emerin Antibody (M00714) overnight at 4°C. Biotinylated goat anti-mouse IgG was used as



secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1021) with DAB as the chromogen.




Figure 4. Flow Cytometry analysis of A431 cells using anti-Emerin antibody (M00714).

Overlay histogram showing A431 cells stained with M00714 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-Emerin Antibody (M00714, 1 µg/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-mouse IgG (BA1126, 5-10 µg/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1 µg/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Emerin EMD Antibody (monoclonal, 5A10)