

## Anti-PKC delta PRKCD Rabbit Monoclonal Antibody

Catalog Number: M00822


### Overview

Product Name	Anti-PKC delta PRKCD Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PKC delta PRKCD Rabbit Monoclonal Antibody catalog # M00822. Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, IHC, ICC, WB
Clonality	Monoclonal IHD-16
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q05655

### Technical Details

Immunogen	A synthesized peptide derived from human PKC delta
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200

## Anti-PKC delta PRKCD Rabbit Monoclonal Antibody (M00822) Images



Western blot analysis of PRKCD using anti-PRKCD antibody (M00822). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HeLa whole cell lysates, Lane 2: human MCF-7 whole cell lysates, Lane 3: rat brain tissue lysates, Lane 4: rat lung tissue lysates, Lane 5: mouse brain tissue lysates, Lane 6: mouse lung tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PRKCD antigen affinity purified monoclonal antibody (Catalog # M00822) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PRKCD at approximately 78 kDa. The expected band size for PRKCD is at 78 kDa.

### 1 Publications Citing This Product

1. PubMed ID: 33745958, Shi Y, Jiang R, Qin X, Gao A, Hou X, Chen L, Xu X, Guo Y, Chai L, Zhao L, Du X, Wu F. Up-regulation of nPKC contributes to proliferation of mice pulmonary artery smooth muscle cells in hypoxia-induced pulmonary hypertension. *Eur J Pharmacol.* 2021 Mar 18:174046. doi: 10.1016/j.ejphar.2021.174046. Epub ahead of print. PMID: 33745958.

Visit [bosterbio.com/anti-pkc-delta-rabbit-monoclonal-antibody-m00822-boster.html](https://www.bosterbio.com/anti-pkc-delta-rabbit-monoclonal-antibody-m00822-boster.html) to see all 1 publications.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PKC delta PRKCD Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.