

## Anti-Cystatin C CST3 Rabbit Monoclonal Antibody

Catalog Number: M00961


### Overview

Product Name	Anti-Cystatin C CST3 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Cystatin C CST3 Rabbit Monoclonal Antibody catalog # M00961. Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Monoclonal IEE-3
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P01034


### Technical Details

Immunogen	A synthesized peptide derived from human Cystatin C
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200

## Anti-Cystatin C CST3 Rabbit Monoclonal Antibody (M00961) Images



Western blot analysis of Cystatin C/CST3 using anti-Cystatin C/CST3 antibody (M00961). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: rat testis tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Cystatin C/CST3 antigen affinity purified monoclonal antibody (M00961) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Cystatin C/CST3 at approximately 16 kDa. The expected band size for Cystatin C/CST3 is at 16 kDa.



Western blot analysis of Cystatin C expression in HeLa cell lysate.

## 1 Publications Citing This Product

1. PubMed ID: 29883558 , BAPTA-AM Nanoparticle for the Curing of Acute Kidney Injury Induced by Ischemia/Reperfusion

Visit [bosterbio.com/anti-cystatin-c-rabbit-monoclonal-antibody-m00961-boster.html](http://bosterbio.com/anti-cystatin-c-rabbit-monoclonal-antibody-m00961-boster.html) to see all 1 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cystatin C CST3 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.