

Anti-EDNRB Antibody Picoband® (monoclonal, 15C3)

Catalog Number: M01041-1

About EDNRB

Endothelin receptor type B, also known as ETB is a protein that in humans is encoded by the EDNRB gene. The protein encoded by this gene is a G protein-coupled receptor which activates a phosphatidylinositol-calcium second messenger system. Its ligand, endothelin, consists of a family of three potent vasoactive peptides: ET1, ET2, and ET3. Studies suggest that the multigenic disorder, Hirschsprung disease type 2, is due to mutations in the endothelin receptor type B gene. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Overview


Product Name	Anti-EDNRB Antibody Picoband® (monoclonal, 15C3)
Reactive Species	Human
Description	Boster Bio Anti-EDNRB Antibody Picoband® (monoclonal, 15C3) catalog # M01041-1. Tested in WB applications. This antibody reacts with Human. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	WB
Clonality	Monoclonal 15C3
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P24530

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human EDNRB, different from the related mouse and rat sequences by one amino acid.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml

Anti-EDNRB Antibody Picoband® (monoclonal, 15C3) (M01041-1) Images



Western blot analysis of EDNRB using anti-EDNRB antibody (M01041-1). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: human placenta tissue lysates, Lane 2: human CACO-2 whole cell lysates, Lane 3: human SW620 whole cell lysates, Lane 4: human HepG2 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-EDNRB antigen affinity purified monoclonal antibody (Catalog # M01041-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for EDNRB at approximately 45KD. The expected band size for EDNRB is at 45KD.

1 Publications Citing This Product

1. PubMed ID: 10.1080/03079457.2013.821568, Expression of endothelin-1 and its receptors in the lungs of broiler chickens exposed to high-altitude hypoxia

Visit bosterbio.com/anti-ednrb-picoband-trade-antibody-monoclonal-m01041-1-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-EDNRB Antibody (monoclonal, 15C3)

For Research Use Only. Not for use in diagnostic procedures.