

Anti-PDPK1/Pdk 1 Rabbit Monoclonal Antibody

Catalog Number: M01159


Overview

Product Name	Anti-PDPK1/Pdk 1 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-PDPK1/Pdk 1 Rabbit Monoclonal Antibody catalog # M01159. Tested in WB, IHC, ICC/IF applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, IHC, ICC, WB
Clonality	Monoclonal HOO-16
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O15530


Technical Details

Immunogen	A synthesized peptide derived from human PDPK1
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200


Anti-PDPK1/Pdk 1 Rabbit Monoclonal Antibody (M01159) Images



Western blot analysis of PDPK1 expression in HEK293 cell lysate.




Western blot analysis of PDPK1 using anti-PDPK1 antibody (M01159). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human 293T whole cell lysates, Lane 2: human HepG2 whole cell lysates, Lane 3: human MCF-7 whole cell lysates, Lane 4: human SIHA whole cell lysates, Lane 5: rat brain tissue lysates, Lane 6: mouse brain tissue lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PDPK1 antigen affinity purified monoclonal antibody (M01159) at 1:1000 overnight at 4°C, then washed with TBS-10% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for PDPK1 at approximately 63 kDa. The expected band size for PDPK1 is at 63 kDa.



Immunohistochemical analysis of paraffin-embedded human kidney, using PDPK1 Antibody.

Immunofluorescent analysis of PC-12 cells, using PDPK1 Antibody.



Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-PDPK1/Pdk 1 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.