

Anti-Bak BAK1 Rabbit Monoclonal Antibody

Catalog Number: M01163


Overview

Product Name	Anti-Bak BAK1 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-Bak BAK1 Rabbit Monoclonal Antibody catalog # M01163. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal CHI-2
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. *This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q16611


Technical Details

Immunogen	A synthesized peptide derived from human Bak
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5mg/ml
Purification	Affinity-chromatography
Suggested Dilutions	WB 1:500-2000 IHC 1:50-200 ICC/IF 1:50-200 IP 1:20 FC 1:20


Anti-Bak BAK1 Rabbit Monoclonal Antibody (M01163) Images



Western blot analysis of Bak using anti-Bak antibody (M01163). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human HeLa whole cell lysates, Lane 2: human 293T whole cell lysates, Lane 3: human Jurkat whole cell lysates, Lane 4: human Daudi whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Bak antigen affinity purified monoclonal antibody (Catalog # M01163) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Bak at approximately 23-25 kDa. The expected band size for Bak is at 23 kDa.




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Immunohistochemical analysis of paraffin-embedded mouse liver, using Bak Antibody.

Immunofluorescent analysis of HeLa cells, using Bak Antibody .



2 Publications Citing This Product

1. PubMed ID: 15248898, BAK overexpression mediates p53-independent apoptosis inducing effects on human gastric cancer cells
2. PubMed ID: 21573215, Minicircle-oriP-IFN?: A Novel Targeted Gene Therapeutic System for EBV Positive Human Nasopharyngeal Carcinoma

Visit bosterbio.com/anti-bak-rabbit-monoclonal-antibody-m01163-boster.html to see all 2 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Bak BAK1 Rabbit Monoclonal Antibody

For Research Use Only. Not for use in diagnostic procedures.