

Anti-STMN1 Antibody (C-Term)

Catalog Number: M01194-2

About STMN1

Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser- 16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear (By similarity).


Overview

Product Name	Anti-STMN1 Antibody (C-Term)
Reactive Species	Human, Rat
Description	Boster Bio Anti-STMN1 Antibody (C-Term) (Catalog # M01194-2). Tested in WB, Flow Cytometry application(s). This antibody reacts with Human, Rat.
Application	Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.
Storage Instructions	Maintain refrigerated at 2-8°C for up to 2 weeks. For long-term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P16949

Technical Details

Immunogen	This STMN1 antibody is generated from a rabbit immunized with a KLH conjugated synthetic peptide between 106-140 amino acids from human STMN1.
Predicted Reactive Species	Bovine, Monkey, Pig
Isotype	Rabbit IgG
Purification	This antibody is purified through a protein A column, followed by peptide affinity purification.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB: 1:2000 FC: 1:25

Anti-STMN1 Antibody (C-Term) (M01194-2) Images



All lanes : Anti-STMN1 Antibody (C-Term) at 1:2000 dilution

Lane 1: HT-29 whole cell lysate

Lane 2: U-2OS whole cell lysate

Lane 3: PC-12 whole cell lysate

Lane 4: HeLa whole cell lysate


Lysates/proteins at 20 µg per lane.

Secondary

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution.

Predicted band size : 17 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing U-2 OS cells stained with M01194-2 (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (M01194-2, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1 µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-STMN1 Antibody (C-Term)