

## Anti-TCF7 Antibody (Monoclonal, 31T28)

Catalog Number: M01315

### About TCF7

Transcription factor 7 is a protein that in humans is encoded by the TCF7 gene. It is mapped to 5q31.1. This gene encodes a member of the T-cell factor/lymphoid enhancer-binding factor family of high mobility group (HMG) box transcriptional activators. This gene is expressed predominantly in T-cells and plays a critical role in natural killer cell and innate lymphoid cell development. The encoded protein forms a complex with beta-catenin and activates transcription through a Wnt/beta-catenin signaling pathway. Mice with a knockout of this gene are viable and fertile, but display a block in T-lymphocyte differentiation. Alternative splicing results in multiple transcript variants. Naturally-occurring isoforms lacking the N-terminal beta-catenin interaction domain may act as dominant negative regulators of Wnt signaling.

### Overview

Product Name	Anti-TCF7 Antibody (Monoclonal, 31T28)
Reactive Species	Human
Description	Boster Bio Anti-TCF7 Antibody (Monoclonal, 31T28) catalog # M01315. Tested in WB, IHC, IP applications. This antibody reacts with Human.
Application	IP, IHC, WB
Clonality	Monoclonal 31T28
Formulation	Rabbit IgG in stabilizing components, phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. This antibody is supplied in a stabilized formulation. Compatibility with conjugation reactions depends on the chemistry of the conjugation method used. For conjugation methods that are not compatible with the stabilizing components present in this formulation, a carrier-free antibody format is required.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P36402


### Technical Details

Immunogen	Recombinant protein within human TCF7 aa 3-298.
Form	Liquid
Concentration	500 ug/ml
Purification	Protein A affinity purified.

Suggested Dilutions

Western blot, 1:500-2000  
Immunohistochemistry, 1:50-200  
ImmunoPrecipitation, 1:50

## Anti-TCF7 Antibody (Monoclonal, 31T28) (M01315) Images



Western blot analysis of TCF7 using anti-TCF7 antibody (M01315). Electrophoresis was performed on a 10% SDS-PAGE gel at 80V (Stacking gel) / 120V (Resolving gel) for 2 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human Jurkat whole cell lysates, Lane 2: human Caco-2 whole cell lysates, Lane 3: human HepG2 whole cell lysates, Lane 4: human SH-SY5Y whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-TCF7 antigen affinity purified monoclonal antibody (M01315) at 1:500 overnight at 4°C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an ECL Plus Western Blotting Substrate (Catalog # AR1196-200) with Tanon 5200 system. A specific band was detected for TCF7 at approximately 42 kDa. The expected band size for TCF7 is at 42 kDa.

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TCF7 Antibody (Monoclonal, 31T28)

For Research Use Only. Not for use in diagnostic procedures.