

Anti-Cytokeratin 8 KRT8 Antibody Picoband® (monoclonal, 3G9)

Catalog Number: M01421-3

About KRT8

Keratin, type II cytoskeletal 8, also known as cytokeratin-8 (CK-8) or keratin-8 (K8) is a keratin protein that is encoded in humans by the KRT8 gene. This gene is a member of the type II keratin family clustered on the long arm of chromosome 12. Type I and type II keratins heteropolymerize to form intermediate-sized filaments in the cytoplasm of epithelial cells. The product of this gene typically dimerizes with keratin 18 to form an intermediate filament in simple single-layered epithelial cells. This protein plays a role in maintaining cellular structural integrity and also functions in signal transduction and cellular differentiation. Mutations in this gene cause cryptogenic cirrhosis. Alternatively spliced transcript variants have been found for this gene.

Overview


Product Name	Anti-Cytokeratin 8 KRT8 Antibody Picoband® (monoclonal, 3G9)
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Cytokeratin 8 KRT8 Antibody Picoband® (monoclonal, 3G9) catalog # M01421-3. Tested in Flow Cytometry, IF, IHC, IHC-F, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Application	Flow Cytometry, IF, IHC, IHC-F, ICC, WB
Clonality	Monoclonal 3G9
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P05787

Technical Details

Immunogen	E.coli-derived human Cytokeratin 8 recombinant protein (Position: D107-K325). Human Cytokeratin 8 shares 95.4% and 94.5% amino acid (aa) sequence identity with mouse and rat Cytokeratin 8, respectively.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(P), IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.

Isotype	Mouse IgG2b
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Western blot, 0.1-0.5ug/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml Immunohistochemistry (Frozen Section), 0.5-1ug/ml Immunofluorescence, 2ug/ml Immunocytochemistry/Immunofluorescence, 2ug/ml Flow Cytometry (Fixed), 1-3ug/1x10 ⁶ cells

Anti-Cytokeratin 8 KRT8 Antibody Picoband® (monoclonal, 3G9) (M01421-3) Images



1. Flow Cytometry analysis of A549 cells using anti-Cytokeratin 8 antibody (M01421-3).

Overlay histogram showing A549 cells stained with M01421-3 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Cytokeratin 8 Antibody (M01421-3, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 2. Western blot analysis of Cytokeratin 8 using anti-Cytokeratin 8 antibody (M01421-3).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human placenta tissue lysates;
Lane 2: human Caco-2 whole cell lysates;
Lane 3: human A549 whole cell lysates;
Lane 4: human A431 whole cell lysates;
Lane 5: human Hela whole cell lysates;
Lane 6: human K562 whole cell lysates;
Lane 7: human HEK293 whole cell lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-Cytokeratin 8 antigen affinity purified monoclonal antibody (Catalog # M01421-3) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for Cytokeratin 8 at approximately 54KD. The expected band size for Cytokeratin 8 is at 54KD.




Figure 3. IHC analysis of Cytokeratin 8 using anti-Cytokeratin 8 antibody (M01421-3).

Cytokeratin 8 was detected in paraffin-embedded section of human rectal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml mouse anti-Cytokeratin 8 Antibody (M01421-3) overnight at 4°C. Peroxidase Conjugated Goat Anti-Mouse IgG was used as secondary antibody and incubated for 30 minutes at

37°C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit (Catalog # SV0001) with DAB as the chromogen.




Figure 4. IF analysis of Cytokeratin 8 using anti-Cytokeratin 8 antibody (M01421-3).

Cytokeratin 8 was detected in paraffin-embedded section of human intestinal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/mL mouse anti-Cytokeratin 8 Antibody (M01421-3) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 5. IF analysis of Cytokeratin 8 using anti-Cytokeratin 8 antibody (M01421-3).

Cytokeratin 8 was detected in paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/mL mouse anti-Cytokeratin 8 Antibody (M01421-3) overnight at 4°C. DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. IF analysis of Cytokeratin 8 using anti-Cytokeratin 8 antibody (M01421-3).

Cytokeratin 8 was detected in immunocytochemical section of U2OS cell. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 2ug/ml mouse anti-Cytokeratin 8 Antibody (M01421-3) overnight at 4°C. Biotin Conjugated Goat anti-Mouse IgG (BA1001) was used as secondary antibody and incubated for 30 minutes at 37°C. The section was developed using Cy3 Conjugated Avidin (BA1037). Visualize using a fluorescence microscope and filter sets appropriate for the label used.

1 Publications Citing This Product

1. PubMed ID: 33636397, Chen H,Kang J,Zhang F,Yan T,Fan W,He H,Huang F.SIRT4 regulates rat dental papilla cell differentiation by promoting mitochondrial functions.Int J Biochem Cell Biol.2021 Feb 23:105962.doi:10.1016/j.biocel.2021.105962.Epub ahead of print.PMID:33636397.

Visit [bosterbio.com/anti-krt8-picoband-trade-antibody-monoclonal-m01421-3-boster.html](https://www.bosterbio.com/anti-krt8-picoband-trade-antibody-monoclonal-m01421-3-boster.html) to see all 1 publications.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Cytokeratin 8 KRT8 Antibody (monoclonal, 3G9)