

# **Anti-PRMT4 CARM1 Monoclonal Antibody**

Catalog Number: M01486-1

### **About CARM1**

TRPC6, also known as TRP6, short transient receptor potential channel 6 and transient receptor potential cation channel subfamily C member 6, is thought to form a receptor-activated non-selective calcium permeant cation channel. TRPC6 is probably operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G-protein coupled receptors. It is activated by diacylglycerol (DAG) in a membrane-delimited fashion, independently of protein kinase C and may not to be activated by intracellular calcium store depletion. Defects in this gene are a cause of focal segmental glomerulosclerosis (FSGS). Expression of this protein has been reported in tissues such as placenta, lung, spleen, ovary, small intestine, and renal podocytes. Immunohistochemistry studies using polyclonal antibodies to this target have shown moderate to strong staining in cell types such as neurons, breast, respiratory, squamous and prostate epithelium, epidermis, placental trophoblasts, dendritic cells, and subsets of immune cells, and faint to moderate staining of adrenal, colon, ganglion cells, hepatocytes, heart, and testis.

#### Overview

Product Name	Anti-PRMT4 CARM1 Monoclonal Antibody
Reactive Species	Human, Monkey, Rat
Description	Boster Bio Anti-PRMT4 CARM1 Monoclonal Antibody catalog # M01486-1. Tested in ELISA, Flow Cytometry, IF, IHC, WB applications. This antibody reacts with Human, Monkey, Rat.
Application	ELISA, Flow Cytometry, IF, IHC, WB
Clonality	Monoclonal 3H2
Formulation	Ascitic fluid containing 0.03% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	Q86X55

#### **Technical Details**

Immunogen	Purified recombinant fragment of human PRMT4 expressed in E. Coli.
Predicted Reactive Species	Chimpanzee
Cross Reactivity	No cross reactivity with other proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml









Purification	Affinity purification
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  WB 1:500-1:2000  IHC 1:200-1:1000  FC 1:200-1:400  ELISA 1:10000




## Anti-PRMT4 CARM1 Monoclonal Antibody (M01486-1) Images




Flow cytometric analysis of Lovo cells using PRMT4 monoclonal antibody (green) and negative control (purple).



Immunofluorescence analysis of Hela cells using PRMT4 monoclonal antibody (green). Red: Actin filaments have been labeled with Alexa Fluor-555 phalloidin.



Immunohistochemistry analysis of paraffin-embedded breast cancer tissues (left) and ovarian cancer tissues (right) with DAB staining using PRMT4 monoclonal antibody.



Western blot analysis using PRMT4 monoclonal antibody against MCF-7 (1), HeLa (2), NIH/3T3 (3), HL-60 (4), LNcap (5), Jurkat (6), PC-3 (7), Cos7 (8), and PC-12 (9) cell lysate.

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.